

RESEARCH ARTICLE

Environmental Mapping of *Paracoccidioides* spp. in Brazil Reveals New Clues into Genetic Diversity, Biogeography and Wild Host Association

Thales Domingos Arantes^{1,2*}, Raquel Cordeiro Theodoro^{2,3}, Marcus de Melo Teixeira^{4,5}, Sandra de Moraes Gimenes Bosco¹, Eduardo Bagagli¹

1 Departamento de Microbiologia e Imunologia, Instituto de Biociências de Botucatu, Universidade Estadual Paulista/UNESP, Botucatu, Brazil, **2** Instituto de Medicina Tropical—IMT/RN, Universidade Federal do Rio Grande do Norte/UFRN, Natal, Brazil, **3** Departamento de Biologia Celular e Genética, Centro de Biociências, Universidade Federal do Rio Grande do Norte/UFRN, Natal, Brazil, **4** Departamento de Biologia Celular, Universidade de Brasília—UnB, Brasília, Brazil, **5** Northern Arizona Center for Valley Fever Research, Translational Genomics Research Institute—Tgen North, Flagstaff, Arizona, United States of America

* smah05@hotmail.com, smah055@yahoo.com.br



OPEN ACCESS

Citation: Arantes TD, Theodoro RC, Teixeira MdM, Bosco SdMG, Bagagli E (2016) Environmental Mapping of *Paracoccidioides* spp. in Brazil Reveals New Clues into Genetic Diversity, Biogeography and Wild Host Association. PLoS Negl Trop Dis 10(4): e0004606. doi:10.1371/journal.pntd.0004606

Editor: Todd Reynolds, University of Tennessee, UNITED STATES

Received: December 2, 2015

Accepted: March 13, 2016

Published: April 5, 2016

Copyright: © 2016 Arantes et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP grant number 2012/03233-3 and 2012/14047-6) and by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Background

Paracoccidioides brasiliensis and *Paracoccidioides lutzii* are the etiological agents of Paracoccidioidomycosis (PCM), and are easily isolated from human patients. However, due to human migration and a long latency period, clinical isolates do not reflect the spatial distribution of these pathogens. Molecular detection of *P. brasiliensis* and *P. lutzii* from soil, as well as their isolation from wild animals such as armadillos, are important for monitoring their environmental and geographical distribution. This study aimed to detect and, for the first time, evaluate the genetic diversity of *P. brasiliensis* and *P. lutzii* for Paracoccidioidomycosis in endemic and non-endemic areas of the environment, by using Nested PCR and *in situ* hybridization techniques.

Methods/Principal Findings

Aerosol (n = 16) and soil (n = 34) samples from armadillo burrows, as well as armadillos (n = 7) were collected in endemic and non-endemic areas of PCM in the Southeastern, Midwestern and Northern regions of Brazil. Both *P. brasiliensis* and *P. lutzii* were detected in soil (67.5%) and aerosols (81%) by PCR of *Internal Transcribed Spacer* (ITS) region (60%), and also by *in situ* hybridization (83%). Fungal isolation from armadillo tissues was not possible. Sequences from both species of *P. brasiliensis* and *P. lutzii* were detected in all regions. In addition, we identified genetic *Paracoccidioides* variants in soil and aerosol samples which have never been reported before in clinical or armadillo samples, suggesting greater genetic variability in the environment than in vertebrate hosts.

Competing Interests: The authors have declared that no competing interests exist.

Conclusions/Significance

Data may reflect the actual occurrence of *Paracoccidioides* species in their saprobic habitat, despite their absence/non-detection in seven armadillos evaluated in regions with high prevalence of PCM infection by *P. lutzii*. These results may indicate a possible ecological difference between *P. brasiliensis* and *P. lutzii* concerning their wild hosts.

Author Summary

Paracoccidioides brasiliensis and *Paracoccidioides lutzii* are the fungal species responsible for one of the most important mycoses of Latin America, Paracoccidioidomycosis (PCM). These fungi can grow in soil from forests, deforested areas, sugarcane, coffee, and rice plantations, as well as pasturelands, and they are strongly associated to armadillo burrows, which can explain their frequent isolation from this mammal's tissues. The environmental detection of these pathogens in endemic and non-endemic areas of PCM is important for mapping risk areas, as well as for understanding the infection ability and clinical manifestations of these fungi. These pieces of information are not provided by isolates obtained from human patients, because these fungi have a long latency period and the human host can migrate, leading to a misinterpretation of the actual geographic distribution of these pathogens. By using two different molecular methodologies (Nested PCR and in situ fluorescence), we detected both species of *P. brasiliensis* and *P. lutzii* in soil and in aerosol samples, even in areas where PCM is only associated to one of these two species. These data might indicate different habitat maintenance strategies between the species, which means that the infection ability may change according to the climatic and soil conditions. Despite contributing new information about the ecology of these important fungal pathogens, our molecular approach for the environmental detection of *Paracoccidioides* species may also be applied for their detection and differentiation in clinical samples, improving the diagnosis of this important systemic mycosis.

Introduction

The study of biological and ecological aspects of *Paracoccidioides brasiliensis* [1,2] and *Paracoccidioides lutzii* [3] has been developed by several research groups in recent years. In particular, investigators have been seeking to isolate and/or detect those pathogens in clinical and environmental samples in order to obtain more in-depth data on the ecological factors that determine their geographical distribution. Both species cause paracoccidioidomycosis (PCM), the most prevalent systemic mycosis in Latin America, which can be acquired by non-immunocompromised hosts inhabiting and/or working mostly in endemic rural areas in South and Central Americas [4,5]. The disease is acquired by soil aerosolization and inhaling of infectious particles of the fungi [5]. It is known that *Paracoccidioides* spp. has its habitat (physical and geographical distribution site) located in the soil, but its ecological niche (sum of all interactions of the microorganism with the biotic and abiotic factors of the environment) has not been properly determined, demanding more environmental studies [4–6].

Studies on *Paracoccidioides* species distribution have been focused mainly on its isolation from human patients and scarcely from environment. Wild and domestic animals have been addressed due to the difficulties to retrieve the fungus from its habitat in laboratory conditions. The few cases of environmental pathogen isolation were from soil, foliage, dog food, bats and

penguin feces, almost serendipitously, with little or no repeatability [4,7–10]. The frequent isolation of *Paracoccidioides brasiliensis* from armadillos, an animal whose home range is very limited, makes this mammal an excellent environmental source for mapping the geographic distribution of the fungus [5,11].

The difficulties faced by environmental studies limits the understanding about the ecology and the real distribution of this genus in endemic and non-endemic areas of PCM. For instance, the majority of the *Paracoccidioides* isolates used for molecular typing are from clinical specimens, which can be influenced by factors such as human migration and the long latency period of this mycosis, which may be greater than one or two decades [12,13], so that it is very hard to specify the exact local of infection and the occurrence of each cryptic species in the endemic and non-endemic regions where the patients are from. Identification of risk areas associated with the different species and/or genotypes can also contribute to a better understanding of *Paracoccidioides* biogeography and also to help the clinical procedures for PCM diagnosis and treatment [11,14]. However, little has been done given its methodological difficulties. In this study, a new approach based on an *in situ* hybridization technique with species-specific DNA probes, as well as the previously established molecular technique of Nested PCR [15,16], were carried out in order to detect the pathogen in different environmental samples, such as soil, aerosol and armadillo tissues. In this study, we expand the collection sites to include new areas of Southeastern, Midwestern and Northern Brazil (prevalent areas of *P. lutzii* infection). The main objective was to search for new ecological and biogeographical information about these fungi, in order to have a better profile of *P. brasiliensis* and *P. lutzii* distribution and dynamics in nature.

Materials and Methods

Ethics statement

Capturing of the armadillos was authorized by SISBIO-IBAMA for all of the Brazilian territory (IBAMA—30585–1 and IBAMA—37333–2), and the Animal Ethics Committee from Biosciences Institute of Botucatu—UNESP (protocol number 528) also approved the procedures.

Selection of the study areas and environmental samples collection

Environmental samples from Rondônia (RO), Goiás (GO) and Minas Gerais (MG) states of Brazil, which have been poorly or never sampled, were selected for the molecular detection of *Paracoccidioides* spp. [17–19]. Samples were obtained by collecting aerosol and soil from armadillo burrows, as well as the animal specimen (*Dasypus novemcinctus*). The distribution of collection areas along the Brazilian map are shown in Fig 1, highlighting the assessed municipalities: Monte-Negro (RO), Santo Antônio de Goiás and Guarani de Goiás (GO) and Campina Verde (MG). Georeferenced sites of the armadillo burrows in the collection areas are listed in Table 1.

Collection of soil and armadillo specimen samples

About 50g of soil was collected from the armadillo burrows by using an iron spatula. The samples were carefully collected so that the burrows were not destroyed. The soil was placed in 50 mL sterile universal bottles, sealed, identified, stored at room temperature, for up to fifteen days, and processed for DNA extraction, accordingly with our previously experience [15,16]. During the collection procedure, the instruments were decontaminated with 70% v/v ethanol solutions in order to avoid cross contamination of samples from one location to another [15,16]. For soil samplings, the minimum established number was 10 samples for burrows and




Fig 1. Collection areas of environmental samples (soil, aerosol and animals) in Brazil. The evaluated states (RO, MG and GO) are highlighted in different colors and the municipalities of sampling in different traces indicated with arrows.

doi:10.1371/journal.pntd.0004606.g001

trails at each site studied. In addition to the collected samples, a soil called "Dark Earth" from the occidental Amazon region (which is an anthropogenic soil and rich in organic matter) was kindly provided by the Center for Nuclear Energy Research Group in Agriculture (CENA)—USP, under the responsibility of PhD. Siu Mui Tsai. These soil samples were processed with the same methods described above (storage and DNA extraction).

For the capture of armadillos in the field, track traps were used and we were assisted by local hunters in order to identify areas with high animal activity. Track traps were placed in the armadillo's trails next to those burrows for which recent animal activity had been detected. In addition to the track traps, active capture took place in the evening, the period of the animal's highest activity.

Captured animals were placed in containers with fresh and dry straw to better accommodate them during transportation until the euthanasia procedure in the laboratory. The number of animals varied according to the season (rain or dry), as well as to the difficulties for finding specimens and transporting them to the laboratory. The euthanasia for the evaluated animals in this study was performed by subcutaneous administration of Zoletyl 50 (0.2 mL/kg/IV, Virbac), followed by cardiac puncture and administration of potassium iodine to ensure the animal's death. Spleen, liver and mesenteric lymph nodes were removed and placed in sterile

Table 1. Georeferenced sites for the field work (soil, aerosols and armadillo collection).

ID	Description/Location	(S) Reference	(WO) Reference
WP01	Trail/SAG	16°29'52,7"	49°17'25,0"
WP02	Burrow/SAG	16°29'53,4"	49°17'25,9"
WP03	Burrow/SAG	16°29'10,3"	49°17'44,1"
WP04	Burrow/SAG	16°29'09,7"	49°17'44,0"
WP05	Burrow/SAG	16°29'37,5"	49°17'27,3"
WP06	Burrow/GDG	13°56'53,9"	46°29'14,6"
WP07	Burrow/GDG	13°51'42,6"	46°31'05,2"
WP08	Burrow/GDG	13°51'43,1"	46°31'05,4"
WP09	Burrow/GDG	13°51'43,3"	46°31'05,2"
WP10	Burrow/GDG	13°51'43,9"	46°31'04,1"
WP11	Embrapa/SAG	16°30'18,1"	49°16'52,0"
WP12	House/Posse-GO	14°05'12,0"	46°21'47,2"
WP13	House/GDG	14°51'40,8"	46°48'10,5"
WP14	House/CV	19°39'48,9"	49°22'07,7"
WP15	Burrow/CV	19°40'00,6"	49°22'03,4"
WP16	Burrow/CV	19°39'59,8"	49°22'01,5"
WP17	Burrow/CV	19°39'59,1"	49°22'02,4"
WP18	Burrow/CV	19°39'59,5"	49°22'02,4"
WP19	Burrow/CV	19°39'54,1"	49°21'54,8"
WP20	Burrow/CV	19°39'57,9"	49°21'48,5"
WP21	Burrow/CV	19°39'57,4"	49°21'47,9"
WP22	Burrow/CV	19°39'57,5"	49°21'50,6"
WP23	Burrow/MN	10°18'46,3"	63°16'57,0"
WP24	Burrow/MN	10°18'56,2"	63°16'56,0"
WP25	Burrow/MN	10°17'23,3"	63°16'33,3"
WP26	Burrow/MN	10°17'19,5"	63°16'57,1"
WP27	Burrow/MN	10°14'41,6"	63°18'14,3"
WP28	Burrow/MN	10°14'05,9"	63°18'16,4"
WP29	Burrow/MN	10°09'09,6"	63°16'18,1"
WP30	Burrow/MN	10°09'09,1"	63°16'17,8"

Legend: Santo Antônio de Goiás (SAG); Guarani de Goiás (GDG); Campina Verde (CV) and Monte Negro (MN); (S)—South; (WO)—East-West; For Armadillo: the WayPoints (WP) 13, 14 and 23 represent areas around the collection sites for captured animals in RO, MG and GO states.

doi:10.1371/journal.pntd.0004606.t001

plates and soaked in alcohol 70% v/v for a brief cleaning followed by saline solution (0,9% w/v). Small fragments (1-2mm) of the organs were then placed in a Mycosel Agar culture medium (Difco) supplemented with 50 µg/mL Gentamicin and incubated at 35°C during 45 days [11,16]. After the evaluation, these plates were properly sterilized and discarded.

Aerosol samples

Air sampler model Cyclone 251 BC, developed by the Centers for Diseases Control—CDC (Morgantown, WV, USA) and certified by the National Institute for Occupational Safety and Health (NIOSH) coupled to vacuum pumps type 224-44XR Model SKC Universal Pumps was used for air sampling [20]. An air sampler was placed next to the armadillo burrows as well as in the areas where the animals had recently removed the soil searching for food. The vacuum pump has a rechargeable battery life up to 24 hours of operation at maximum flow, thereby

facilitating the procedures for collection in remote locations without any power sources. Aerosol and soil samples were collected in the states of GO and MG during the dry/warm season, and in RO state during the cold/rainy season. Each aerosol collection was performed in a minimum period of 60 minutes with a flow rate of 3500 mL/min. At least four samples were collected at each site.

DNA extraction and PCR/nested PCR reactions

Each soil sample was subjected to total DNA extraction in triplicate, using the commercial kit PowerSoil DNA Isolation Kit—MO BIO Laboratories, Inc. Total DNA was resuspended in 100 µl of Nuclease free water and quantified in NanoVue spectrophotometer equipment (GE Healthcare). Fifty microliters of each one of the three replicates were mixed in a single 1.5 mL tube and concentrated to a final volume of 30µl in a concentrator (Eppendorf) and then quantified again to confirm the new concentration of each sample. The aerosol samples were directly used for PCR without any previous DNA extraction by washing the tubes from cyclonic sampler with the PCR reaction mix.

The PCRs (Polymerase Chain Reaction) were carried out with ITS-4/5 primers for rRNA universal fungal region ITS1-5.8S-ITS2 (*Internal Transcribed spacer*) [21]. A Nested PCR was performed with the product of the first amplification using specific primers for the *Paracoccidioides* genus, annealing in the ITS-1 and ITS-2 regions, named PbITS-E (5'GAGCTTTGAC GTCTGAGACC3') and PbITS-T (5'GTATCCCTACCTGATCCGAG3') [16]. Both PCR mixes were prepared using 12.5 µl of Nuclease Free Water (Sigma), 0.5 µl of 0.2 mM dNTP mix, 5.0 µl of 5X GC buffer, 2.5 µl of 30% DMSO, 0.625 µl of each primer at 20 µM and 0.25 µl of 1000 units/µl Taq Phusion DNA Polymerase (ThermoFisher) for each reaction with 23.0 µl of PCR mix for 2.0 µl from a DNA soil sample of approximately 15 ng/µl. Twenty five microliters of PCR reaction was performed for aerosol samples, these mixes were prepared from 100.0 µl of reaction mix used for washing the aerosol collection tubes, so that four PCRs were carried out for each aerosol sample.

The thermal cycling conditions for the first PCR were: an initial denaturation at 98°C for 30 seconds and 39 cycles of denaturation at 98°C for 10 seconds, followed by an annealing step at 55°C for 45 seconds, and extension at 72°C for 45 seconds, after that a final extension at 72°C for 10 minutes was applied. For aerosol samples, the first step of denaturation was longer (5 minutes) than the one applied to the DNA soil samples, in order to break the spores and other fungal structures, releasing the genetic material in the PCR mix. For Nested PCR, the annealing step was adjusted to 58°C.

After the PCR and Nested PCR reactions, PCR products were analyzed by electrophoresis in a 1.5% w/v agarose gel. The bands around 450bp (*Paracoccidioides* spp.) or that were best highlighted in the gel were cut out, purified by using the commercial Kit (GE illustrates GFX PCR DNA and Gel Band Purification) and quantified as described above. Purified samples were sent to the Laboratory for Molecular Diagnosis of the Department of Microbiology and Immunology (UNESP, Botucatu/SP-Brazil) for automatic capillary sequencing in ABI 3500 DNA Analyzer (Applied Biosystems) equipment. For low concentration of PCR products, a new amplification reaction with the PbITS-E/T primers were performed (double PCR), followed by its purification and sequencing.

The obtained sequences were aligned to the reference sequences with the help of the MEGA 6.0 program [22] and compared to an online database (GenBank) [23], to verify their identity and phylogenetic clusterization with other deposited ITS sequences from *P. brasiliensis* and *P. lutzii*.

Phylogenetic analysis

The initial sequences obtained from environmental amplicons were checked using the Sequencing Analysis software from ABI 3500 DNA Analyzer (Applied Biosystems) in order to improve the quality of sequences. For phylogenetic analysis and comparison of other *Paracoccidioides* DNAs, we used the deposited sequences under the following access numbers: EU870314; EU870315; EU870316; AY631235; EU118561; EU118560; EU118548; EU118554; EU118553; EU118549; EU118546; EU118547; EU118545; EU118543; EU118542 for *P. brasiliensis* and EU870298; EU870303; EU870306; EU870309; EU870310; EU870311; AF092903; EU870299 for *P. lutzii*. Only the sequences, herein obtained from environmental samples, presenting $\geq 97\%$ of similarity to *Paracoccidioides* spp. sequences in *GenBank* (blastn analysis) [24] were considered for phylogenetic analysis in MEGA 6.0 software. Sequences previously generated from soil and aerosol samples from the Southeast part of Brazil [16] were included in this dataset, access numbers KP636439 to KP636474 (S1 Table). In addition, clinical strain sequences from *GenBank*, representing *P. lutzii* (Pb01—EU870297), *P. brasiliensis* species complex S1 (Pb18—AF322389), PS2 (Pb3—EU870315), and PS3 (AY631237) were included into the final dataset. Sequences were aligned with the ClustalW [25] algorithm implemented in the Bioedit software [26]. Retrieved alignments were manually inspected in order to avoid mispaired bases. Neighbor-Joining [27] and Maximum Likelihood trees were inferred in the MEGA 6 software [22] using the Jukes-Cantor [28] nucleotide substitution model. One thousand bootstrap replicates [29] were used to estimate the monophyletic clades support, and values were displayed next to the branches.

Detection and differentiation of *P. brasiliensis* complex and *P. lutzii* species by *in situ* hybridization in environmental aerosol samples

Probes used in this study were commercially synthesized targeting the rRNA region of *P. brasiliensis* and *P. lutzii*, specifically for the ITS-1-5.8S-ITS-2 region [14,30,31]. Sixty ITS sequences from different *Paracoccidioides* isolates were aligned in order to select conserved regions within species, exclusive and different between *P. brasiliensis* and *P. lutzii* (Fig 2). The probes were differentially labeled on their 5' end, with Horseradish Peroxidase (HRP) for *P. brasiliensis*, and Texas Red for *P. lutzii*, (Fig 2).

The use of this approach for detection of *Paracoccidioides* spp. as well as differentiation between the *P. brasiliensis* complex and *P. lutzii* species was previously standardized [32]. For those methods, each probe was tested against other Ajellomycetaceae fungi (*Histoplasma capsulatum*) and other clinically relevant ascomycetes (*Aspergillus flavus*; *Aspergillus fumigatus*; *Trichophyton mentagrophytes*), as negative controls for both probes (used in FISH and TSA--FISH techniques). Twelve aerosol samples were collected in armadillo burrows for *in situ* hybridization method: four from the state of GO, four from MG and four from RO.

The detection of *Paracoccidioides* spp. with DNA probes by the TSA-FISH method in aerosol samples was applied for greater sediment volume in the cyclone sampler tubes (stages). These samples from both stages of the cyclone sampler were fixed with 1.5 mL of 4% Paraformaldehyde plus 0.1 M Phosphate solution buffer. Series of 50%, 80% and 100% ethanol solutions were used to remove cell fixation solution and to dehydrate the cells, so that they have the ability to absorb the probes to be used in the hybridization step. After dehydration, 10 mL of pre-hybridization buffer [2.0 mL of ultra-pure water; 4.0 mL of 40% Formamide; 1.8 mL of 5 M NaCl; 200 μ l of 1 M Tris (pH 7.5); 100 μ l of 1% SDS; 2 mL of 10% Buffer Blocking Agent] were added to the samples for stabilization and improvement of their permeability. After this first step, cells were hybridized with probes at a final concentration of 50 ng/ μ l in hybridization buffer. After 16–17 hours of incubation at 42°C, the slides with fungal controls were washed




Fig 2. The ITS region with the location of oligo probes designed for the detection and differentiation of *P. brasiliensis* and *P. lutzii*, labeled with HRP and Texas Red, respectively.

doi:10.1371/journal.pntd.0004606.g002

with 50 mL of Washing Buffer [47.54 mL of ultra-pure water; 460 µl of 5 M NaCl, 500 µl of 0.5M EDTA, 500 µl of 1% SDS and 1 mL of 1 M Tris (pH 7.5)] for removal of non-specific binding probes. After washing, the slides were stabilized with 250 mL of TNT buffer [217.315 mL of ultra-pure water; 25 mL of 1 M Tris (pH 7.5); 7.5 mL of 5M NaCl and 0.185 mL of Tween 20].

After equilibrating and washing the slides with TNT buffer, 30 µl of TSA solution (TSA Plus PerkinElmer Kit) were added to each slide and incubated for 30 minutes in a humid dark chamber at room temperature. The slides were washed again, dried at room temperature, prepared with the addition of 4',6-Diamidino-2-phenylindole dihydrochloride (DAPI) and covered with a cover slip to be observed under a fluorescence microscope. These slides were divided in two groups of four slides each: one was tested against *P. brasiliensis* probe and the other against *P. lutzii* probe. Two spare slides were used as controls during the hybridization phase for each of the two methods and probes used in this study.

Results

Animals evaluated in the collected areas

Seven armadillos were captured and evaluated, three from GO, three from RO and one from MG states. Information about the gender, weight and cultivated organ fragments are listed in [Table 2](#).

Table 2. Armadillos (*Dasypus novemcinctus*), and their respective organ fragments, evaluated in the Brazilian states of Goiás, Minas Gerais and Rondônia.

ID Animal	Sex	Weight	Spleen (Plates)	Liver (Plates)	L. M. (Plates)	Total of Plates	Number of Fragments
AGO1	Male	5.5 kg	12	66	06	84	1,950
AGO2	Male	4.3kg	09	53	02	64	1,966
AGO3	Male	4.5kg	21	44	03	68	2,548
ARO1	Female	5.0kg	12	20	05	37	1,050
ARO2	Female	4.3kg	12	20	05	37	1,060
ARO3	Male	1.8kg	10	20	06	36	1,000
AMG1	Male	4.5kg	15	48	07	70	2,450
Total of Organ Fragments							12,024

Legend: AGO—Armadillos from Goiás; ARO—Armadillos from Rondônia; AMG—Armadillos from Minas Gerais; L.M.—mesenteric lymphnodes; Plates—Total number of culture plates assessed for each organ; Number of Fragments—total number of fragments from each organ that were cultured in Mycosel agar for each animal evaluated.

doi:10.1371/journal.pntd.0004606.t002

After 45 days of incubation, each plate was evaluated for fungal growth similar to *Paracoccidioides* spp. by micro-morphological analysis. All the colonies presented morphological structures of bacteria, despite the addition of antibiotics to the culture medium and no fungal structures were identified. After 45 days, the plates were then considered negative for *Paracoccidioides* spp. growth.

Nested PCR, phylogenetic and haplotype analysis of soil and aerosol samples

Forty-four soil samples from armadillo burrows were obtained; 12 in GO, 10 in MG and 22 in RO. Twenty-eight aerosol samples from armadillo burrows were obtained; 10 in GO, 10 in MG and 08 in RO, 16 of these samples were set aside for molecular detection by Nested PCR and the rest of the 12 aerosol samples were used for *in situ* hybridization techniques. The sampled burrows in the areas of RO, GO and MG states were mostly located in deforested areas of pastures or in some riparian forest sites. Positive Nested PCR amplification for ITS region of *Paracoccidioides* spp. was observed in 67.5% of soil samples and in 81% of aerosol samples. No amplicons were observed for “Dark earth” soil samples. When compared to the GenBank database, sequences revealed SNPs specific for *P. lutzii* and/or *P. brasiliensis* in all the positive soil samples from RO, GO and MG. *P. lutzii* was found in aerosol samples in MG and GO states, while *P. brasiliensis* was only detected in GO (Table 3).

Phylogenetic analyses were carried out using a total of 36 sequences obtained from the current study and 11 sequences collected from the GenBank were added to the final dataset. The majority of the soil and aerosol samples clustered within *P. lutzii* species (27 out of 36 –Fig 3A). *P. lutzii* was detected in GO, RO and MG states, as well as in São Paulo (SP) as previously reported [16]. All these *P. lutzii* sequences were displayed in a single haplotype together with the referenced clinical strain Pb01 (Fig 3B). Solely 4 samples (AR_GO1, AR_GO19, AR_GO11 and AR_GO2D) were clustered within *P. brasiliensis*, all obtained from aerosol samples collected in GO state (Fig 3A). However, only the AR_GO1 and AR_GO19 samples are clustered with the clinical strains Pb3, Pb18 and ATCC60855 in unique haplotype representing clinical isolates from the *P. brasiliensis* species complex S1, PS2 and PS3 (Fig 3B).

The phylogenetic distribution and haplotypic network analysis revealed higher genetic variation in the environmental samples than reported so far for *Paracoccidioides* clinical samples. The samples AR_GO11 and AR_GO2D are disposed to polytomic branches within *P. brasiliensis* and constitute single haplotypes in the network (Fig 3). Moreover, two high supported clades were observed as being closely related to *P. lutzii* clinical/environmental samples (Soil clades). The soil clade I is composed of the soil samples SO_GO10 and SO_GO19 from GO state, while the soil clade II is composed of soil samples SO_RO11 and SO_RO12 from RO state. In addition, the AR_MG16 sample collected from aerosol in MG also appears to be a genetic variant from *P. lutzii*, fallen into a paraphyletic branch in the tree (Fig 3A).

Detection and differentiation of *P. brasiliensis* and *P. lutzii* in aerosol samples by *in situ* hybridization

The detection and differentiation of *Paracoccidioides* spp. in aerosol samples was performed after validation of FISH and TSA-FISH in *P. brasiliensis* and *P. lutzii* in culture cells. Specificity and sensitivity control tests were applied for validating the positive detection in environmental samples [32]. The probes tested (conjugated with HRP and Texas Red) did not hybridize with the negative controls. Both *P. brasiliensis* and *P. lutzii* cells under culture conditions show specific nuclear staining, which merge with DAPI.

Table 3. Detection of *Paracoccidioides* spp. by Nested PCR and *in situ* hybridization techniques, both in soil and aerosol samples in the states of Goiás, Minas Gerais and Rondônia, Brazil.

Location	Nested PCR					<i>in situ</i> hybridization			
	Soil		Total # of collected samples	Aerosol		Total # of collected samples/ PCR Reactions*	Aerosol		Total # of samples
	Pb	PI		Pb	PI		Pb	PI	
Goiás	0	9 (75%)	12	4 (16.5%)	0	6/24	4 (100%)	2 (50%)	4
Minas Gerais	0	0	10	0	9 (37.5%)	6/24	2 (50%)	0	4
Rondônia	0	14 (64%)	22	0	0	4/16	1 (25%)	1 (25%)	4
Positivity: 67.5%			Positivity: 81%			Positivity: 83%			

Legend: *Pb*—*Paracoccidioides brasiliensis*; *PI*—*Paracoccidioides lutzii*;

*—Total # of collected samples/ PCR Reactions shows the number of total samples collected; for each sample, the PCR mix was divided in 4 separated reactions. The positivity of Nested PCR for the aerosol samples is based on the number of positive amplification out of the total of PCR reactions.

doi:10.1371/journal.pntd.0004606.t003

For the *in situ* hybridization, the aerosol samples with higher sediment in the tubes were obtained in dry areas (GO and MG) where the burrow soil was easily aerosolized. For the RO state samples, which were obtained during an intense rainy season, the pellets were less visible to the naked eye.

From the 72 slides prepared for the *in situ* hybridization, positive hybridization occurred in 36 (50% of the slides), revealing that ten out of 12 air samples from the armadillo burrows (83%) were positive for *Paracoccidioides* spp. (Table 3). The *in situ* hybridization experiments using specific probes for *P. brasiliensis* and *P. lutzii* revealed the presence of both species in GO and RO states (Fig 4A, 4B, 4E and 4F), and only *P. brasiliensis* species in MG state (Fig 4).

Fig 5 summarizes the current data on *Paracoccidioides* spp. detection in soil and aerosol samples by the different methodologies applied so far, including the current results and those previously obtained [16].

Discussion

Soil and aerosol samples have shown to be excellent sources for mapping the fungus in endemic areas of PCM by molecular methods, since they are easy to obtain and handle in a laboratory, therefore corroborating previous studies of our group [15,16]. Both soil and aerosol samples were positive for the environmental detection of *Paracoccidioides* spp. DNA in the sampled areas, revealing the ubiquitous distribution of these pathogens in the Brazilian territory. Aerosol samples were collected in a smaller number compared with the soil samples, due to the methodological difficulties in the field and weather conditions of each sampled location. The soil collection methodology is faster and easier to perform than the aerosol sampling, which requires more field effort to become representative. On the other hand, the aerosol sampling mimics the fungal dispersion mechanism by which rural workers become infected with the mycelia and/or conidia particles. Currently, soil from endemic or non-endemic areas for PCM may contain fungal cells with infectious potential for human and/or animal population, although the course of infections can vary according to the biotic and abiotic factors of the environment. It has been stated that agricultural activities are the activities that most favor PCM infection in humans [34,35]. The aerosol samples are also methodologically simple to work with in the laboratory because DNA extraction is not required.




Fig 3. A) Molecular Phylogenetic Analysis by of ITS locus revealed by Maximum Likelihood and Neighbor Joining methods, using the Jukes-Cantor model parameters with range correction. Replication percentages on the tree are grouped in the bootstrap test (1000 replicates) and shown next to the branches. The sequences related to environmental samples are identified by acronyms SO_GO (Soil of Goiás) and AR_GO (Aerosol Goiás), AR_MG (Aerosol Minas Gerais) and SO_RO (Soil of Rondônia). **B)** Median-joining network showing the unique haplotypes of the Soil Clades I and II. Circles are proportional to haplotype frequency and numbers of mutations are represented by black dots. Red circles represent hypothetical missing intermediates (median vectors).

doi:10.1371/journal.pntd.0004606.g003




Fig 4. Fungal structures (400X) visualized by FISH and TSA-FISH techniques for aerosol samples and controls. **A:** aerosol samples from Goiás with DAPI. **B:** aerosol samples from Goiás with *P. brasiliensis* probe. **C:** aerosol sample from Rondônia, with DAPI. **D:** aerosol samples from Rondônia with *P. brasiliensis* probes. **E:** aerosol sample from Goiás with DAPI. **F:** aerosol sample from Goiás with *P. lutzii* probe. **G and I:** *Histoplasma capsulatum* with DAPI. **H:** *Histoplasma capsulatum* with *P. brasiliensis* probe (specificity control). **J:** *Histoplasma capsulatum* with *P. lutzii* probe (specificity control). **K:** isolate Pb01 (*P. lutzii*) with *P. lutzii* probe (positive control). **L:** isolate T16B1 (*P. brasiliensis*) with *P. brasiliensis* probe. The probe for *P. brasiliensis* is conjugated with Horseradish Peroxidase/HRP and *P. lutzii* probe is labeled with Texas Red/txr, and genetic material is labeled with DAPI.

doi:10.1371/journal.pntd.0004606.g004




Fig 5. Brazilian sites and biomes where environmental detection (from soil, aerosol and armadillos) of *Paracoccidioides* spp. has been carried out. The collection areas encompass the states of Minas Gerais (MG), Goiás (GO) and Rondônia (RO) states, to São Paulo (SP) and Pará (PA). Circles outside of the map indicate the percentage of positivity in each area for *P. brasiliensis* (green) and *P. lutzii* (red) by *in situ* hybridization for aerosol samples. The circles outside of the map in black and grey indicate the percentage of positivity in each area for *P. brasiliensis* and *P. lutzii* by Nested PCR for soil and aerosol samples. The white circle indicates the negative detection in the evaluated areas (N.D. = Not detected). The armadillos indicate the number of collected animals and the positivity for isolation of *Paracoccidioides brasiliensis* (Pb) in each location of our study (in RO, GO and MG) and from previous works in SP [16] and PA [33]. The triangles (green), squares (yellow) and circles (red) show the distribution of the clinical isolates for cryptic species (S1, PS2 and *P. lutzii*) in previous studies [14,31].

doi:10.1371/journal.pntd.0004606.g005

Molecular detection in soil samples was positive in two of the sampled areas (RO and GO), but it was negative in MG. The molecular detection of *Paracoccidioides* in RO soil showed higher amplification rate as visualized in the agarose gel electrophoresis (S1 Fig), probably because of the rainy conditions during the collection, which according to Barrozo et al., 2009 and 2010 [34,35], may favor fungus maintenance in soil, as well as its dispersion, causing increased incidence of PCM. Positive samples were also obtained from northeastern Goiás, a warm and dry area, which might indicate some fungal resistance to adverse conditions.

This work presented the first standardization of *in situ* hybridization techniques for environmental search of *Paracoccidioides* spp., (HRP-probe/TSA-FISH for *P. brasiliensis* and Texas Red-probe/FISH for *P. lutzii*), making its detection possible in aerosol samples from the three locations studied (RO, MG and GO). This new approach showed sensitivity and specificity rates similar to the well-established Nested PCR technique. The advantage of *in situ*

hybridization is the visualization of infective fungal structures of *Paracoccidioides* spp. directly in the environmental samples.

The detection rate of *Paracoccidioides* spp. in soil and aerosol samples in GO, MG and RO was lower than the detection rate in endemic areas of PCM. In most cases, the fungus was detected in locations whose air humidity and temperature conditions were very similar to those found in endemic areas of PCM. However, new distribution nuances of *P. brasiliensis* and *P. lutzii* were revealed in our study, including a remarkable resistance to adverse environmental conditions, so that the spatial distribution of the *Paracoccidioides* species may be larger than previously defined based mainly on clinical isolates [30,31,36].

Both growth and dispersion of this pathogen seems to be greatly influenced by the climate. While high moisture levels increase fungal growth and maintenance in soil, a brief drought period dries the most superficial layer of the soil, making the dispersion of aerosols (mycelia particles/conidia and other microorganisms) easy and intense [37]. This was already observed in *Coccidioides* spp. [38,39] and could explain the negative detection of *Paracoccidioides* spp. in aerosol samples from RO, which faced one of the most severe rainy periods of the last few years [40]. This explains the higher positive molecular detection of *Paracoccidioides* spp. by Nested PCR in soil samples than in aerosol samples from RO, where the collection was carried out during the raining season. Therefore, different from soil samples, which directly demonstrate the presence of the fungus, and aerosol samples also reflect the spread of the fungus in the environment, and therefore its infective potential is extremely important in the epidemiological study of *Paracoccidioides* spp., corroborating the growth and blow theory which reflects the crescent number of cases after rainy seasons in endemic areas [34,35] and the great recent number of new PCM cases in areas of North Brazil, as RO state [19].

In RO and GO, the detection of *Paracoccidioides* spp. was positive mostly in deforested pastures and in some riparian forest sites. In fact, such areas with increased agricultural activity present the greatest incidence of PCM cases in these states [17–19]. Deforestation of preserved areas exposes the soil and naturally or deliberately changes its chemical conformation, which can favor the infection of rural workers or other people living in these areas, leading to the emergence of PCM. Our studies indicate that *P. lutzii* and *P. brasiliensis* is often found in soil and aerosol samples in all four of the sampled regions. For this reason, we hypothesized that the geographic distribution of PCM caused by different *Paracoccidioides* species may be associated to the capacity of each fungal species to produce infective propagule in the current environmental conditions, which includes soil management for local agriculture activity (sugarcane, coffee and cattle breeding). For instance, despite the environmental molecular detection of *P. brasiliensis* in Goiás and *P. lutzii* in Minas Gerais, the majority of PCM cases in these states are caused by *P. lutzii* and *P. brasiliensis*, respectively.

Considering the difficulty for animal capture and transportation, a reasonable number of armadillos for each sampled area were obtained. All armadillos evaluated in this study were negative for culture isolation of *Paracoccidioides* spp. However, molecular detection in soil and aerosol samples indicated the presence of *Paracoccidioides* spp. in these areas. This negative result for the non-endemic areas, where these animals were collected (northeastern GO and MG), may point to the relationship of environmental conditions and the possibility of infection and/or disease in these animals, and probably in humans too. On the other hand, the negative isolation in armadillos from the endemic area (RO) is more intriguing. In this area, most of the sequences detected showed high similarity to *P. lutzii* species (in all positive soil samples by Nested PCR, and in 50% of the positive aerosol samples by *in situ* hybridization). This may indicate that the relationship between armadillo and *P. lutzii* is different from the well-known interaction between armadillo and *P. brasiliensis*, which could have resulted in a speciation process (*P. brasiliensis* complex X *P. lutzii*) driven by hosts. *P. brasiliensis* S1 and PS2 species

(São Paulo state/Brazil) and PS3 (Colombia) being highly recovered from the armadillos, and *D. novemcinctus* and *Cabassous centralis* [11,41] in endemic areas, while no *P. lutzii* has been isolated from these animals yet.

According to the phylogenetic analysis, there was a prevalence of sequences belonging to the *P. lutzii* species in all evaluated areas, which reflect the distribution pattern of clinical isolates observed in recent works [14,16,18]. Despite the negative isolation of *Paracoccidioides* spp. from armadillos, data from molecular detection of these pathogens in soil and/or aerosol samples may be useful for delineating the geographic distribution of *Paracoccidioides* spp.

Herein for the first time, environmental genetic variants were reported for *Paracoccidioides* genus (Fig 3). Phylogenetic and haplotype data revealed the presence of two well-supported clades in environmental soil samples, one from GO (Soil Clade I) and the other from RO (Soil Clade II). The sample AR_MG16 collected from aerosol in MG also appears to be a genetic variant from *P. lutzii*, fallen into a paraphyletic branch in a tree. It's worth noting that all environmental samples that clustered apart from the clinical referenced strains of both *P. lutzii* and *P. brasiliensis* are displayed in single haplotypes, reinforcing a higher diversity of the *Paracoccidioides* spp. in the environment than in human hosts. This discrepancy shows that part of the environmental genotypes of *Paracoccidioides* spp. may not be able to infect and/or cause PCM.

It is interesting to note that although Central-Western Brazil presents a prevalence of PCM caused by *P. lutzii*, *P. brasiliensis* was also detected in these areas in this work and in previous studies [30,31,42]. This observation could indicate different patterns of sporulation depending on soil constitution and weather, so that the conidia production/release of *P. brasiliensis* and *P. lutzii* could be different in distinct areas, explaining the current distribution pattern of PCM caused by both species. Previous studies have already pointed out differences in sporulation (concerning conidia amount and morphology) among the different cryptic species: S1 isolates produce a higher conidia amount compared to other *P. brasiliensis* and *P. lutzii* species. This observation may explain the higher isolation of S1 species (from humans and armadillos) in endemic areas for PCM caused by *P. brasiliensis* [42,43].

Molecular epidemiology studies of fungal pathogens are extremely important, since the genetic variability may reflect the existence of cryptic species, can have a geographic pattern and also be related to different clinical manifestations and antifungal drug response. For that reason, many efforts have been made to study the geographic limits of the different genotypes of some fungal pathogens, such as *Histoplasma capsulatum*, a complex of at least seven distinct clades, whose variability has been addressed by MLST (Multi-locus sequencing type), as well as ITS and PRP8 intein genes as molecular markers [44–46]. Cryptic speciation event has also been recently described for another Ajellomycetacea member, *Blastomyces dermatitidis*, which is now considered two species, *B. dermatitidis* and *Blastomyces gilchristii*. Like *P. brasiliensis* and *P. lutzii*, these *Blastomyces* species seem to be sympatric in some of their distribution ranges [47].

Despite that ITS region is widely used as fungal barcoding [48], most studies on cryptic speciation use the multi-locus sequencing approach. That is because ITS is, in some cases, useless to distinguish among very close cryptic species, such as those from the *P. brasiliensis* complex [31]. However *P. brasiliensis* and *P. lutzii* are clearly distinguished by ITS sequencing. Thus, our findings in fact confirm the coexistence of the *P. lutzii* and *P. brasiliensis* species (S1 and PS2 species) in the environment, even though the potential for human infection seems to be different for both species, depending on the region.

This study presents important data regarding the eco-epidemiology of *Paracoccidioides* species, as well as their actual distribution over the Brazilian map. In addition, new environmental genetic variants of these pathogens should notify us about the actual scenario of the *Paracoccidioides* species diversity. Our results also pointed out the possible differential pathogen versus

wild host interaction among *Paracoccidioides* species, generating new hypotheses and new work issues to be tested and studied by the scientific community. Thus, further studies based on data interacting with clinical and ecological aspects should be conducted in order to create a more reliable distribution map of this important systemic mycosis and its etiological agents. In addition, higher throughput sequencing methods should be addressed in order to get a better resolution of *Paracoccidioides* species diversity and distribution.

Supporting Information

S1 Table. Sequences obtained from environmental amplicon with ITS1;5.8S;ITS2 partial sequences deposited in the Genbank platform from the NCBI database.

(DOC)

S1 Fig. Imagens of agarose gel electrophoresis by molecular detection of ITS1;5.8S;ITS2 of *Paracoccidioides* spp. in soil and aerosol samples of RO, GO and MG states of Brazil.

Images (A, B, C, D, E and F).

(DOC)

Acknowledgments

We thank PhD. Siu Mui Tsai and her students for the DNA soil samples from the Amazon. We also thank the employers and farmers from the collection areas who helped us collect the environmental samples and animals. We thank the Ph.D Brett Green from CDC (Centers for Diseases Control and Prevention, Morgantown, WV, USA) for providing the cyclonic sampler used in this study.

Author Contributions

Conceived and designed the experiments: TDA RCT EB. Performed the experiments: TDA MdMT SdMGB. Analyzed the data: TDA RCT MdMT. Contributed reagents/materials/analysis tools: TDA RCT MdMT SdMGB EB. Wrote the paper: TDA RCT MdMT SdMGB EB.

References

1. Splendore A. Blastomycoses Americanas. *Rev Bras Med.* 1910; 24: 153–7.
2. Almeida FP. Estudos comparativos do granuloma coccidióico nos Estados Unidos e no Brasil. Novo gênero para o parasito brasileiro. *An Fac Med Univ Sao Paulo.* 1930; 5: 125–41.
3. Teixeira MM, Theodoro RC, Nino-Veja G, Bagagli E, Felipe MS. *Paracoccidioides* species complex: ecology, phylogeny, sexual reproduction, and virulence. *PLoS Pathog.* 2014; 10(10): e1004397. doi: [10.1371/journal.ppat.1004397](https://doi.org/10.1371/journal.ppat.1004397) PMID: [25357210](https://pubmed.ncbi.nlm.nih.gov/25357210/)
4. Franco M, Bagagli E, Scapolio S, Lacaz CS. A critical analysis of isolation of *Paracoccidioides brasiliensis* from soil. *Med Mycol.* 2000; 38 (3): 185–191 doi: [10.1080/mmy.38.3.185.191](https://doi.org/10.1080/mmy.38.3.185.191) PMID: [10892985](https://pubmed.ncbi.nlm.nih.gov/10892985/)
5. Bagagli E, Theodoro RC, Bosco SM, McEwen JG. *Paracoccidioides brasiliensis*: phylogenetic and ecological aspects. *Mycopathologia.* 2008; 165 (4–5): 197–207. PMID: [18777629](https://pubmed.ncbi.nlm.nih.gov/18777629/)
6. Restrepo A. The ecology of *Paracoccidioides brasiliensis*: a puzzle still unsolved. *Sabouraudia.* 1985; 23(5): 323–34. PMID: [3906945](https://pubmed.ncbi.nlm.nih.gov/3906945/)
7. Shome SK, Batista AC. Occurrence of *Paracoccidioides brasiliensis* In the soil of Recife (Brazil). *Rev. Fac Med Univ Ceara.* 1963; 3: 90–94.
8. Grose E, Tramsitt JR. *Paracoccidioides brasiliensis* recovered from the intestinal tract of three bats (*Artibeus lituratus*) in Colombia, S.A. *Sabouraudia.* 1965; 4(2): 124–125. PMID: [5896105](https://pubmed.ncbi.nlm.nih.gov/5896105/)
9. Gezeule E. Aislamiento de *Paracoccidioides* sp. de heces de pinguino de la Antartida. In: San-Blas G. ed. *Proceedings IV Encuentro Internacional sobre Paracoccidioidomicosis.* Caracas, Venezuela: Instituto Venezolano de Investigaciones Cientificas (IVIC). 1989; Abstract B2.

10. Silva-Vergara ML, Martinez R, Chadu A, Madeira M, Freitas-Silva G, Leite Maffei CM. Isolation of *Paracoccidioides brasiliensis* strain from the soil of a coffee plantation in Ibiá, State of Minas Gerais, Brazil. *Med Mycol.* 1998; 36(1): 37–42. PMID: [9776810](#)
11. Bagagli E, Franco M, Bosco SM, Hebeler-Barbosa F, Trinca LA, Montenegro MR. High frequency of *Paracoccidioides brasiliensis* infection in armadillos (*Dasypus novemcinctus*): an ecological study. *Med Mycol.* 2003; 41(3): 217–223. PMID: [12964713](#)
12. Restrepo A. Morphological aspects of *Paracoccidioides brasiliensis* in lymph nodes: implications for the prolonged latency of paracoccidioidomycosis? *Med Mycol.* 2000 Aug; 38(4):317–22. PMID: [10975700](#)
13. Shikanai-Yasuda MA, Telles Filho FQ, Mendes RP, Colombo AL, Moretti ML. Guidelines in paracoccidioidomycosis. *Revista da Sociedade Brasileira de Medicina Tropical* 39(3):297–310, may-jun, 2006. PMID: [16906260](#)
14. Theodoro RC, Teixeira MM, Felipe MS, Paduan Kdos S, Ribolla PM, San-Blas G, et al. *Paracoccidioides* Genus: Species Recognition and Biogeographic Aspects. *PLoS One.* 2012; 7(5): e37694. Epub 2012 May 30. PMID: [22666382](#)
15. Theodoro RC, Candeias JM, Araújo JP Jr, Bosco Sde M, Macoris SA, Padula LO, et al. Molecular detection of *Paracoccidioides brasiliensis* in soil. *Med Mycol.* 2005; 43 (8): 725–729. PMID: [16422303](#)
16. Arantes TD, Theodoro RC, Da Graça Macoris SA, Bagagli E. Detection of *Paracoccidioides* spp. in environmental aerosol samples. *Med Mycol.* 2013; 51(1): 83–92. doi: [10.3109/13693786.2012.698444](#) Epub 2012 Jul 5. PMID: [22762209](#)
17. Ferreira SP, Santos RS, de Paula LB, Reis AAS. Prevalence of paracoccidioidomycosis in patients diagnosed in Araujo Jorge Hospital, Goiânia-Goiás, Brazil. *Rev Univ Vale do Rio Verde.* 2012; 10: 167–177.
18. Gegembauer G, Araujo LM, Pereira EF, Rodrigues AM, Paniago AM, Hahn RC, et al. Serology of paracoccidioidomycosis due to *Paracoccidioides lutzii*. *PLoS Negl Trop Dis.* 2014; 8(7): e2986. doi: [10.1371/journal.pntd.0002986](#) eCollection 2014. PMID: [25032829](#)
19. Vieira Gde D, Alves Tda C, Lima SM, Camargo LM, Sousa CM. Paracoccidioidomycosis in a western Brazilian Amazon State: clinical-epidemiologic profile and spatial distribution of the disease. *Rev Soc Bras Med Trop.* 2014; 47(1): 63–8. doi: [10.1590/0037-8682-0225-2013](#) PMID: [24603739](#)
20. Green BJ, Schmechel D, Tovey ER. Detection of aerosolized *Alternaria alternata* conidia, hyphae, and fragments by using a novel double-immunostaining technique. *Clin Diagn Lab Immunol.* 2005; 12(9): 1114–1116. PMID: [16148180](#)
21. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. *PCR Protocols: A guide to methods and applications.* San Diego: Academic. 1990; 315–322.
22. Tamura K, Stecher G, Peterson D, Filipksi A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol.* 2013; 30(12): 2725–9. doi: [10.1093/molbev/mst197](#) Epub 2013 Oct 16. PMID: [24132122](#)
23. Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. GenBank. *Nucleic Acids Res.* 2016; 44 (D1): D67–72. doi: [10.1093/nar/gkv1276](#) Epub 2015 Nov 20. PMID: [26590407](#)
24. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic Local Alignment Search Tool. *J Mol Biol.* 1990; 215: 403–410. PMID: [2231712](#)
25. Thompson JD, Higgins DG, Gibson TJ. Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 1994; 22(22): 4673–4680. PMID: [7984417](#)
26. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser.* 1999; 41: 95–98.
27. Saitou N, Nei M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 1987; 4: 406–425. PMID: [3447015](#)
28. Jukes TH, Cantor CR. Evolution of protein molecules. In: Munro HN, editor. *Mammalian Protein Metabolism.* 1969; 21–132.
29. Felsenstein J. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution.* 1985; 39: 783–791.
30. Matute DR, McEwen JG, Puccia R, Montes BA, San-Blas G, Bagagli E, et al. Cryptic speciation and recombination in the fungus *Paracoccidioides brasiliensis* as revealed by gene genealogies. *Mol Biol Evol.* 2006; 23(1): 65–73. Epub 2005 Sep 8. PMID: [16151188](#)
31. Teixeira MM, Theodoro RC, de Carvalho MJ, Fernandes L, Paes HC, Hahn RC, et al. Phylogenetic analysis reveals a high level of speciation in the *Paracoccidioides* genus. *Mol Phylogenet Evol.* 2009; 52(2): 273–83. doi: [10.1016/j.ympev.2009.04.005](#) Epub 2009 Apr 17. PMID: [19376249](#)

32. Arantes, TD. Detecção do *Paracoccidioides* spp. em amostras ambientais e diferenciação do complexo *P. brasiliensis* da espécie *P. lutzii* por Nested PCR e hibridização *in situ*. 2015. 94 f. Tese (doutorado)—Universidade Estadual Paulista Júlio de Mesquita Filho, Instituto de Biociências de Botucatu, 2015. Available at: <<http://hdl.handle.net/11449/132750>>.
33. Naiff RD, Ferreira LCL, Barrett TV, Naiff MF, Arias JR. Paracoccidioidomicose enzoótica em tatus (*Dasyus novemcinctus*) no estado do Pará. Rev Inst Med Trop Sao Paulo. 1986; 28: 19–27.
34. Barrozo LV, Mendes RP, Marques SA, Benard G, Silva ME, Bagagli E. Climate and acute/subacute Paracoccidioidomycosis in a hyper-endemic area in Brazil. Int J Epidemiol. 2009; 38(6):1642–9. doi: [10.1093/ije/dyp207](https://doi.org/10.1093/ije/dyp207) Epub 2009 May 11. PMID: [19433518](https://pubmed.ncbi.nlm.nih.gov/19433518/)
35. Barrozo LV, Bernard G, Silva ME, Bagagli E, Marques SA, Mendes RP. First description of a cluster of acute/subacute Paracoccidioidomycosis cases and its association with a climate-anomaly. PLoS Negl Trop Dis. 2010; 4(3): e643. doi: [10.1371/journal.pntd.0000643](https://doi.org/10.1371/journal.pntd.0000643) PMID: [20361032](https://pubmed.ncbi.nlm.nih.gov/20361032/)
36. Carrero LL, Niño-Vega G, Teixeira MM, Carvalho MJ, Soares CM, Pereira M, et al. New *Paracoccidioides brasiliensis* isolate reveals unexpected genomic variability in this human pathogen. Fungal Genet Biol. 2008; 45(5): 605–12. doi: [10.1016/j.fgb.2008.02.002](https://doi.org/10.1016/j.fgb.2008.02.002) Epub 2008 Feb 21. PMID: [18364259](https://pubmed.ncbi.nlm.nih.gov/18364259/)
37. Yamamoto N, Bibby K, Qian J, Hospodsky D, Rismani-Yazdi H, Nazaroff WW, et al. Particle-size distributions and seasonal diversity of allergenic and pathogenic fungi in outdoor air. ISME J. 2012; 6(10): 1801–11. doi: [10.1038/ismej.2012.30](https://doi.org/10.1038/ismej.2012.30) Epub 2012 Apr 5. PMID: [22476354](https://pubmed.ncbi.nlm.nih.gov/22476354/)
38. Tamerius JD, Comrie AC. Coccidioidomycosis Incidence in Arizona Predicted by Seasonal Precipitation. PLoS One. 2011; 6(6): e21009. doi: [10.1371/journal.pone.0021009](https://doi.org/10.1371/journal.pone.0021009) Epub 2011 Jun 20. PMID: [21701590](https://pubmed.ncbi.nlm.nih.gov/21701590/)
39. Comrie AC, Glueck MF. Assessment of climate-coccidioidomycosis model: model sensitivity for assessing climatologic effects on the risk of acquiring coccidioidomycosis. Ann N Y Acad Sci. 2007; 1111: 83–95. Epub 2007 Mar 7. PMID: [17344540](https://pubmed.ncbi.nlm.nih.gov/17344540/)
40. INPE. National Institute of Aerospace Research—Brazil: <http://www.inpe.br/> (research in 08/02/2015—SINDA Integrated System of Environmental Data).
41. Corredor GG, Peralta LA, Castaño JH, Zuluaga JS, Henao B, Arango M, et al. The naked-tailed armadillo *Cabassous centralis* (Miller 1899): a new host to *Paracoccidioides brasiliensis*. Molecular identification of the isolate. Med Mycol. 2005; 43(3): 275–280. PMID: [16010854](https://pubmed.ncbi.nlm.nih.gov/16010854/)
42. Molinari-Madlum EE, Felipe MS, Soares CM. Virulence of *Paracoccidioides brasiliensis* isolates can be correlated to groups defined by random amplified polymorphic DNA analysis. Med Mycol. 1999; 37(4): 269–276. PMID: [10421862](https://pubmed.ncbi.nlm.nih.gov/10421862/)
43. Terçarioli GR, Bagagli E, Reis GM, Theodoro RC, Bosco Sde M, Macoris AS, et al. Ecological study of *Paracoccidioides brasiliensis* In soil: growth ability, conidia production and molecular detection. BMC Microbiol. 2007; 7: 92. PMID: [17953742](https://pubmed.ncbi.nlm.nih.gov/17953742/)
44. Kasuga T, White TJ, Koenig G, McEwen J, Restrepo A, Castañeda E, et al. Phylogeography of the fungal pathogen *Histoplasma capsulatum*. Mol Ecol. 2003 Dec; 12(12): 3383–401. PMID: [14629354](https://pubmed.ncbi.nlm.nih.gov/14629354/)
45. Landaburu F, Cuestas ML, Rubio A, Elías NA, Daneri GL, Veciño C, et al. Genetic diversity of *Histoplasma capsulatum* strains isolated from Argentina based on nucleotide sequence variations in the internal transcribed spacer regions of rDNA. Mycoses. 2014; 57(5): 299–306. doi: [10.1111/myc.12159](https://doi.org/10.1111/myc.12159) Epub 2013 Dec 2. PMID: [24299459](https://pubmed.ncbi.nlm.nih.gov/24299459/)
46. Theodoro RC, Scheel CM, Brandt ME, Kasuga T, Bagagli E. PRP8 intein in cryptic species of *Histoplasma capsulatum*: evolution and phylogeny. Infect Genet Evol. 2013; 18:174–82. doi: [10.1016/j.meegid.2013.05.001](https://doi.org/10.1016/j.meegid.2013.05.001) Epub 2013 May 10. PMID: [23665464](https://pubmed.ncbi.nlm.nih.gov/23665464/)
47. Brown EM, McTaggart LR, Zhang SX, Low DE, Stevens DA, Richardson SE. Phylogenetic analysis reveals a cryptic species *Blastomyces gilchristii*, sp. nov. within the human pathogenic fungus *Blastomyces dermatitidis*. PLoS One. 2013; 8(3): e59237. doi: [10.1371/journal.pone.0059237](https://doi.org/10.1371/journal.pone.0059237) Epub 2013 Mar 22. PMID: [23533607](https://pubmed.ncbi.nlm.nih.gov/23533607/)
48. Schoch CL, Seifertb KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, et al. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. PNAS. 2012; 109(16): 6241–6. doi: [10.1073/pnas.1117018109](https://doi.org/10.1073/pnas.1117018109) Epub 2012 Mar 27. PMID: [22454494](https://pubmed.ncbi.nlm.nih.gov/22454494/)