

Brazilian Journal of Botany



All the contents of this journal, except where otherwise noted, is licensed under a Creative Commons Attribution License. Fonte:
https://www.scielo.br/scielo.php?script=sci_arttext&pid=S0100-84042012000400011&lng=en&nrm=iso&tlang=en. Acesso em: 16 out. 2020.

REFERÊNCIA

SILVA, Juliana Silvestre; FELFILI, Jeanine Maria. Floristic composition of a conservation area in the Federal District of Brazil. **Brazilian Journal of Botany**, São Paulo, v. 35, n. 4, p. 377-384, 2012. DOI: <https://doi.org/10.1590/S0100-84042012000400011>. Disponível em: https://www.scielo.br/scielo.php?script=sci_arttext&pid=S0100-84042012000400011&lng=en&nrm=iso&tlang=en. Acesso em: 16 out. 2020.

Floristic composition of a conservation area in the Federal District of Brazil¹

JULIANA SILVESTRE SILVA^{2,4} and JEANINE MARIA FELFILI^{3†}

(received: March 19, 2012; accepted: August 10, 2012)

ABSTRACT – (Floristic composition of a conservation area in the Federal District of Brazil). *Cerradão* vegetation shares many species with savanna and forest areas and is one of the most vulnerable phytobiognomies in the *Cerrado* (Brazilian savanna) biome. The floristic composition of the Cerradão Biological Reserve was examined between September/2007 and November/2008. A total of 282 species distributed among 194 genera and 75 families were encountered, demonstrating proportions of 0.91 herbaceous species and 0.54 shrub species for each tree species. Fabaceae, Asteraceae, Rubiaceae, Poaceae, Myrtaceae, Malpighiaceae, and Melastomataceae were the most species-rich families. Fully 72.3% of the species of this dystrophic cerradão were shared by cerrado and forest vegetations, while 60.43% were shared by other cerradão sites, although the largest proportion of species (91%) were shared with cerrado sensu stricto. No species was found to be exclusive to this cerradão site, but approximately 95% of all species were native to the *Cerrado* biome.

Key words - APA Gama e Cabeça de Veado, cerradão, flora, species richness

INTRODUCTION

Cerradão is typified by a continuous arboreal layer more than 7 m tall, with 50 to 90% total canopy coverage; the shrub layer is generally dense and composed largely of sclerophyllous species, while there is only a reduced herbaceous layer. Lianas and some climbers are common, but epiphytes are rare (Ribeiro & Walter 2008). Although cerradão constitutes a distinct *cerrado* (Brazilian savanna) phytobiognomy, it does not have a distinctive flora, but rather a mix of floristic elements from cerrado sensu stricto and forests (Felfili 2002).

Very few studies have focused on the herbaceous/subshrub flora of cerradão, as this is a predominantly arboreal phytobiognomy. Additionally, although an extensive survey of the flora of the Gama and Cabeça de Veado Environmental Protection Area – EPA (in portuguese: *Área de Proteção Ambiental* – APA) was undertaken, the Cerradão Biological Reserve (in portuguese, *Reserva Biológica* – REBIO), also located in this APA, has not been examined in any detail. The present study was designed

to contribute to future planning and management of the area and to our general knowledge of the mixed flora of this under-protected phytobiognomy.

MATERIAL AND METHODS

The present study was undertaken in the Cerradão REBIO, a conservation area created by Federal Decree nº 19.213/98 as an Area of Relevant Ecological Interest (in portuguese, *Área de Relevante Interesse Ecológico* – ARIE) and later transformed into a Biological Reserve (REBIO) by Federal Decree nº 31.757/10. This conservation area covers 54 ha (15°51' S and 47°49' W) and encompasses cerradão as well as dense cerrado vegetation. The regional climate is Tropical Altitudinal (Cwa by the Köppen classification system), with average temperatures in the coldest month below 18 °C, and above 22 °C in the warmest month; the average annual rainfall varies from 1,350 mm to 1,450 mm. This REBIO is extremely important in protecting the water resources of the Federal District, as the region supplies approximately 50% of the local aquifer recharge volume (Ibram 2010). The soils of the reserve are red latosols, similar to those found in other cerradão areas of the Gama and Cabeça de Veado EPA (Área de Preservação Ambiental – APA Gama e Cabeça de Veado) in the Federal District (Unesco 2002).

The floristic composition of the cerradão vegetation in the REBIO was examined by undertaking collections every two weeks during the period between September/2007 and November/2008. The cerradão vegetation (which differs from the adjacent dense cerrado vegetation by its more robust phytobiognomy) within the reserve was surveyed by following a zigzag route through its entire area.

1. Part of the Masters thesis of the first author, Programa de Pós Graduação em Botânica, Universidade de Brasília, Brasília, DF, Brazil.

2. Universidade de Brasília, Instituto de Ciências Biológicas, Departamento de Botânica, Caixa Postal 4457, 70910-970 Brasília, DF, Brazil.

3†. Universidade de Brasília, Faculdade de Tecnologia, Departamento de Engenharia Florestal, Laboratório de Manejo Florestal, Caixa Postal 04357, 70919-970 Brasília, DF, Brazil (in memoriam).

4. Corresponding author: julissilva@unb.br

Characteristics such as plant size and color and the presence of odors or latex were noted. Species identifications were carried out by comparisons with herbarium samples deposited in the University of Brasília (UB), Brasília Botanical Garden (HEPH), and IBGE (IBGE) herbaria, as well as by consulting the literature and taxonomists. In addition to the species collected during our survey, the floristic list includes species encountered during a phytosociological study of the arboreal layer of the same area undertaken by Silva (2009). Some of the species collected in the latter study were only encountered in their vegetative phase in the present survey and were not assigned collection numbers.

The data gathered in the present survey were compared to plant lists published by Mendonça et al. (2008) as the vascular flora of the Cerrado biome, by Proença et al. (2001) as the phanerogamic flora of the Federal District, and by Felfili et al. (2004) as the vascular flora of the Gama and Cabeça de Veadó EPA, as well as in posterior publications as the phanerogamic flora of the Federal District series (Assis 2002, Carvalho-Silva & Cavalcanti 2002, Lombardi 2002, Mendonça & Amaral Junior 2002, Ribeiro & Proença 2002, Chacon et al. 2003, Ferreira et al. 2003, Groppo 2003, Lima & Pirani 2003, Pontes & Mello-Silva 2003, Silva & Valente 2003, França 2005, Silva & Barros 2005, Fiaschi 2006, Martins & Filgueiras 2006, Pastore & Cavalcanti 2006, Dittrich et al. 2007, Groppo & Pirani 2007).

The present survey considered: trees belonging to the arboreal layer, shrubs belonging to the shrub-layer, subshrubs, herbaceous plants, and climbers (lianas in general) composing the herbaceous layer, in order to allow comparisons with other floristic surveys, including Proença et al. (2001), Felfili et al. (2004), and Mendonça et al. (2008). Voucher specimens were

deposited in the UB herbarium, and duplicates incorporate into the HEPH and UFMT (Universidade Federal do Mato Grosso) herbaria.

Synonyms were verified by consulting taxonomic revisions of specific groups as well as the databank sites of the Missouri Botanical Garden – W³Tropicos (2008) and the International Plant Names Index – IPNI (2008). The botanical classification system followed the recommendations of the Angiosperm Phylogeny Group II (APG II 2003) for the phanerogams flora, and Smith et al. (2006) for the pteridoflora.

RESULTS AND DISCUSSION

We encountered 282 species belonging to 194 genera and 75 plant families in the Cerradão REBIO, of which 115 (40.78%) were arboreal species, 62 (21.99%) were shrubs, and 105 (37.23%) made up the herbaceous stratum (54 herbs, 31 subshrubs, and 20 climbers) (table 1), resulting in ratios of 0.91 herbaceous species and 0.54 shrub species for each arboreal species. The data of Felfili et al. (2004) for cerradão sites in the Gama and Cabeça de Veadó EPA in the Federal District (DF) yielded portions of 0.67 herbaceous species and 0.41 shrub species for each tree species; Proença et al. (2001) found 0.53 herbaceous plant species and 0.45 shrubs per arboreal species in cerradão sites in the Federal District; while Mendonça et al. (2008) reported portions of 1.05 herbaceous species and 0.61 shrubs for each arboreal layer species in cerradão sites in the Cerrado biome.

Table 1. The vascular flora of *cerradão* vegetation in the “REBIO Cerradão”, Federal District, Brazil. Species identified by family, stratum, and phytophysiognomy according to the literature: Proença et al. 2001, Assis 2002, Carvalho-Silva & Cavalcanti 2002, Lombardi 2002, Mendonça & Amaral Junior 2002, Ribeiro & Proença 2002, Chacon et al. 2003, Ferreira et al. 2003, Groppo 2003, Lima & Pirani 2003, Pontes & Mello-Silva 2003, Silva & Valente 2003, Felfili et al. 2004, França 2005, Silva & Barros 2005, Fiaschi 2006, Martins & Filgueiras 2006, Pastore & Cavalcanti 2006, Dittrich et al. 2007, Groppo & Pirani 2007, Mendonça et al. 2008. (* = exotic/invasive; S = savanna sensu stricto; S(ls) = savanna sensu lato; C = *cerradão*; F = forest; GF = gallery forest; MF = mesophytic/dry forest; -- = absent from the lists; Coll. N. = number of collection of JS Silva).

Family	Species	Stratum	Phytophysiognomy	Coll. N.
Acanthaceae	<i>Justicia lanstyakii</i>	Herb	S(ls), S, GF	16
Alstroemeriaceae	<i>Alstroemeria gardneri</i>	Herb	S	369
Amaranthaceae	<i>Gomphrena agrestis</i>	Herb	S, C	482
	<i>Pfafia denudata</i>	Herb	S	218
Anacardiaceae	<i>Anacardium humile</i>	Shrub	S, C	4
	<i>Astronium fraxinifolium</i>	Tree	S, C, GF, MF	20
Annonaceae	<i>Annona crassiflora</i>	Tree	S, C, GF	236
	<i>A. tomentosa</i>	Shrub	S, C, GF	149
	<i>Duguetia furfuracea</i>	Shrub	S, C, GF	169
	<i>Guatteria sellowiana</i>	Tree	F, GF	--
	<i>Xylopia aromatica</i>	Tree	S, C, GF, MF	69

continue

continuation

Family	Species	Stratum	Phytophysiognomy	Coll. N.
Apocynaceae	<i>X. brasiliensis</i>	Tree	S, C, GF	--
	<i>X. sericea</i>	Tree	C, F, GF, MF	57
	<i>Aspidosperma macrocarpon</i>	Tree	S, C, MF	154
	<i>A. spruceanum</i>	Tree	S(ls), GF, MF	--
	<i>A. subincanum</i>	Tree	F, GF, MF	--
	<i>A. tomentosum</i>	Tree	S, C, GF, MF	501
	<i>Ditassa obcordata</i>	Shrub	S, C, F, GF	447
	<i>Ditassa retusa</i>	Climber	S, GF	614
	<i>Hancornia speciosa</i>	Tree	S, C	72
Araliaceae	<i>Odontadenia lutea</i>	Climber	S, C, GF, MF	209
	<i>Prestonia coalita</i>	Climber	S(ls), F, GF, MF	358
	<i>Ilex concolor</i>	Tree	S, C, GF	7
Arecaceae	<i>Schefflera macrocarpa</i>	Tree	S, C, GF	172
Asteraceae	<i>Allagoptera campestris</i>	Shrub	S, C	203
	<i>Syagrus comosa</i>	Shrub	S, C, GF	356
	<i>S. flexuosa</i>	Shrub	S, C, GF	74
	<i>Achyrocline satureoides</i>	Herb	S(ls), S, GF	547
	<i>Aspilia foliacea</i>	Herb	S	243
	<i>Baccharis dracunculifolia</i>	Subshrub	S(ls), GF	263
	<i>B. salzmannii</i>	Shrub	S(ls)	120
	<i>Campuloclinium megacephalum</i>	Subshrub	S, F, GF	384
	<i>Chromolaena laevigata</i>	Shrub	S, F, GF	405
Balanophoraceae	<i>C. leucocephala</i>	Shrub	S	177
	<i>Echinocoryne holosericea</i>	Herb	S	534
	<i>Elephantopus mollis</i> *	Herb	S, C, F, GF, MF	527
	<i>Emilia fosbergii</i> *	Herb	S(ls)	258
	<i>Eremanthus glomerulatus</i>	Shrub	S, C, GF	159
	<i>Gochnatia floribunda</i>	Shrub	S(ls), S, F	73
	<i>Hoehnephytum trixoides</i>	Subshrub	S	46
	<i>Ictyothere latifolia</i>	Subshrub	S, F	368
	<i>Lepidaploa aurea</i>	Shrub	S, GF	99
	<i>Lessingianthus compactiflorus</i>	Subshrub	S, GF	362
Bignoniaceae	<i>Piptocarpha macropoda</i>	Tree	S, F, GF, MF	115
	<i>P. rotundifolia</i>	Tree	S, C	487
	<i>Sympphyopappus compressus</i>	Subshrub	S, GF	292
	<i>Trichogonia salviaefolia</i>	Herb	S, GF	259
	<i>Tridax procumbens</i> *	Herb	MF	386
	<i>Vernonia rubriflava</i>	Shrub	S, C, GF	87
	<i>Wedelia bishopii</i>	Herb	S	344
	<i>Langsdorffia hypogaea</i>	Herb	C, F, GF	517
	<i>Adenocalymma pedunculatum</i>	Shrub	S	213
	<i>Anemopaegma acutifolium</i>	Subshrub	S	519
Blechnaceae	<i>Cuspidaria sceptrum</i>	Climber	S, C, GF, MF	176
	<i>Fridericia platyphylla</i>	Climber	S, C, GF	381
	<i>Handroanthus ochraceus</i>	Tree	S, C, GF	98
	<i>H. serratifolius</i>	Tree	S(ls), S, F, GF, MF	59
Boraginaceae	<i>Jacaranda ulei</i>	Subshrub	S, GF	103
	<i>Zeyheria montana</i>	Shrub	S, C	118
Bromeliaceae	<i>Blechnum australe</i> subsp. <i>auriculatum</i>	Herb	--	274
	<i>Cordia truncata</i>	Subshrub	S	389
Burseraceae	<i>Bromeliaceae</i> sp.	Herb	--	520
	<i>Protium ovatum</i>	Shrub	S, F, GF, MF	25

continue

continuation

Family	Species	Stratum	Phytophysiognomy	Coll. N.
Caryocaraceae	<i>Caryocar brasiliense</i>	Tree	S, C	95
Celastraceae	<i>Cheiloclinium cognatum</i>	Tree	GF	277
	<i>Plenckia populnea</i>	Tree	S, C, GF	205
	<i>Salacia crassifolia</i>	Shrub	S, GF	70
Chrysobalanaceae	<i>Couepia grandiflora</i>	Tree	S, C, GF, MF	502
Clusiaceae	<i>Kielmeyera abdita</i>	Shrub	S	235
	<i>K. coriacea</i>	Tree	S, C, GF	81
	<i>K. speciosa</i>	Tree	S, F, GF	--
Combretaceae	<i>Terminalia argentea</i>	Tree	S, C, GF	77
	<i>T. fagifolia</i>	Tree	S, C, GF, MF	109
Connaraceae	<i>Connarus suberosus</i>	Tree	S, C	556
Convolvulaceae	<i>Ipomoea procumbens</i>	Climber	S	328
	<i>Merremia contorquens</i>	Climber	S	467
Cucurbitaceae	<i>Cayaponia tayuya</i>	Climber	S, F, GF	343
Cunoniaceae	<i>Lamanonia ternata</i>	Tree	S(ls), C, F, GF	--
Cyperaceae	<i>Cyperus laxus</i>	Herb	S(ls), GF	270
	<i>Rhynchospora consanguinea</i>	Herb	S, GF	347
	<i>R. exaltata</i>	Herb	S, C, F	47
Dichapetalaceae	<i>Tapura amazonica</i>	Tree	GF, MF	603
Dilleniaceae	<i>Davilla elliptica</i>	Shrub	S, GF	155
Dioscoreaceae	<i>Dioscorea dodecaneura</i>	Climber	GF	604
Ebenaceae	<i>Diospyros hispida</i>	Tree	S, C, GF, MF	--
Erythroxylaceae	<i>Erythroxylum daphnites</i>	Tree	S(ls), S, C, GF, MF	--
	<i>E. deciduum</i>	Shrub	S, C, F, GF	607
	<i>E. suberosum</i>	Shrub	S, F	152
Euphorbiaceae	<i>Dalechampia caperonioides</i>	Herb	S	342
	<i>Manihot violacea</i>	Shrub	S	337
	<i>Maprounea guianensis</i>	Tree	S, C, GF, MF	5
Fabaceae	<i>Acosmium dasycarpum</i>	Tree	S, C, GF	326
	<i>Bauhinia cf. dumosa</i>	Shrub	S	425
	<i>B. rufa</i>	Subshrub	S, C, GF	608
	<i>Bowdichia virgiliooides</i>	Tree	S, C, GF, MF	557
	<i>Calliandra dysantha</i>	Subshrub	S(ls), S, C, GF	43
	<i>Centrosema brasiliianum</i>	Climber	S(ls), GF	473
	<i>Chamaecrista conferta</i>	Subshrub	S	85
	<i>Copaifera langsdorffii</i>	Tree	S, C, GF, MF	577
	<i>Crotalaria flavigoma</i>	Herb	S(ls), S, GF	376
	<i>C. cf. grandiflora</i>	Shrub	S(ls), GF	418
	<i>Dalbergia miscolobium</i>	Tree	S, C, GF	233
	<i>Dimorphandra mollis</i>	Tree	S, C, GF	237
	<i>Enterolobium gummiferum</i>	Tree	S, C, GF	--
	<i>Galactia boavista</i>	Herb	S(ls)	214
	<i>G. grewiaeefolia</i>	Subshrub	S, C	372
	<i>Hymenaea stigonocarpa</i>	Tree	S, C, GF	271
	<i>Hymenolobium heringerianum</i>	Tree	F, GF, MF	--
	<i>Machaerium acutifolium</i>	Tree	S, C, GF, MF	60
	<i>M. opacum</i>	Tree	S, C, MF	--
	<i>Mimosa somnians</i>	Subshrub	S(ls), S, C, GF	345
	<i>M. velloziana *</i>	Shrub	S(ls), GF	599
	<i>Periandra mediterranea</i>	Shrub	S, C, GF	91
	<i>Piptadenia gonoacantha</i>	Tree	GF, MF	611
	<i>Plathymenia reticulata</i>	Tree	S, C	452

continue

continuation

Family	Species	Stratum	Phytophysiognomy	Coll. N.
	<i>Platypodium elegans</i>	Tree	S(ls), S, C, GF, MF	192
	<i>Pterodon pubescens</i>	Tree	S, C, GF	112
	<i>Stryphnodendron adstringens</i>	Tree	S, C	490
	<i>Stylosanthes guianensis</i>	Herb	S(ls), S	114
	<i>Tachigali vulgaris</i>	Tree	S, C, GF	--
	<i>Vatairea macrocarpa</i>	Tree	S, C, GF	--
Gentianaceae	<i>Voyria aphylla</i>	Herb	F, GF	399
Hypericaceae	<i>Vismia glaziovii</i>	Tree	C, F, GF	267
Icacinaceae	<i>Emmotum nitens</i>	Tree	S, C, GF	83
Lamiaceae	<i>Aegiphila lhotzkiana</i>	Tree	S, C, GF	244
	<i>Hypenia brachystachys</i>	Subshrub	S(ls), S, C, GF	538
	<i>Hyptis lythroides</i>	Subshrub	S	127
	<i>H. rubiginosa</i>	Subshrub	S	107
	<i>H. villosa</i>	Herb	S	133
Lauraceae	<i>Cassytha filiformis</i> *	Herb	S	364
	<i>Endlicheria paniculata</i>	Tree	GF	254
	<i>Nectandra reticulata</i>	Tree	GF	--
	<i>Ocotea aciphylla</i>	Tree	GF	--
	<i>O. diospyrifolia</i>	Tree	S(ls), S, GF	498
	<i>O. pomaderroides</i>	Tree	S(ls), S, GF	9
	<i>O. spixiana</i>	Tree	S, C, GF	21
Loganiaceae	<i>Antonia ovata</i>	Tree	S, C, GF	497
	<i>Strychnos pseudoquina</i>	Tree	S, C	28
Loranthaceae	<i>Phthirusa ovata</i>	Herb	S, GF, MF	151
	<i>Psittacanthus robustus</i>	Herb	S, GF	226
Lythraceae	<i>Cuphea spermacoce</i>	Subshrub	S, C	242
	<i>Diplusodon virgatus</i>	Shrub	S, C, GF	456
	<i>Lafoensia pacari</i>	Tree	S, C, GF, MF	--
Malpighiaceae	<i>Banisteriopsis anisandra</i>	Climber	S(ls), S, GF	78
	<i>B. argyrophylla</i>	Climber	S(ls), S, C, GF, MF	359
	<i>B. megaphylla</i>	Shrub	S, GF	469
	<i>B. stellaris</i>	Climber	S, C, GF	331
	<i>Byrsonima crassifolia</i>	Tree	S, C	79
	<i>B. intermedia</i>	Tree	S(ls), GF	183
	<i>B. laxiflora</i>	Tree	S(ls), S, GF, MF	150
	<i>B. rotunda</i>	Tree	--	221
	<i>B. sericea</i>	Tree	S(ls), S, GF	--
	<i>B. verbascifolia</i>	Tree	S, GF	145
	<i>Heteropterys byrsonimifolia</i>	Tree	S	24
	<i>H. pteropetala</i>	Shrub	S, C, F, GF	330
	<i>Janusia</i> sp.	Climber	--	137
	<i>Peixotoa goiana</i>	Shrub	S	478
	<i>P. reticulata</i>	Shrub	S	543
Malvaceae	<i>Eriotheca pubescens</i>	Tree	S(ls), S, C, F, GF	--
	<i>Pavonia rosa-campestris</i>	Herb	S, C	117
	<i>Pseudobombax longiflorum</i>	Tree	S, CS, GF, MF	561
	<i>Sida linifolia</i> *	Herb	S, GF	625
	<i>Waltheria indica</i> *	Shrub	S, GF	525
Melastomataceae	<i>Leandra aurea</i>	Shrub	S(ls), GF	336
	<i>Miconia albicans</i>	Tree	S, C, GF	2
	<i>M. burchellii</i>	Tree	S, C, GF, MF	6
	<i>M. cubatanensis</i>	Tree	S, C, GF, MF	529

continue

continuation

Family	Species	Stratum	Phytophysiognomy	Coll. N.
	<i>M. cuspidata</i>	Tree	S(ls), S, C, GF, MF	562
	<i>M. fallax</i>	Shrub	S, C	33
	<i>M. ferruginata</i>	Tree	S, C, GF	8
	<i>M. pepericarpa</i>	Tree	S, GF	14
	<i>M. pohliana</i>	Tree	S, C, GF	558
	<i>Miconia sellowiana</i>	Tree	S, C, GF, MF	573
	<i>Ossaea congestiflora</i>	Shrub	S, F	276
	<i>Tibouchina stenocarpa</i>	Tree	S, C, F, GF	426
Menispermaceae	<i>Cissampelos pareira</i> *	Climber	GF	609
Moraceae	<i>Brosimum gaudichaudii</i>	Shrub	S, C	66
Myristicaceae	<i>Virola sebifera</i>	Tree	S(ls), C, GF, MF	26
Myrsinaceae	<i>Cybianthus densiflorus</i>	Subshrub	S, F, GF	126
	<i>C. detergens</i>	Subshrub	S(ls), S, C, GF	247
	<i>Myrsine coriacea</i>	Tree	S(ls), C, GF, MF	147
	<i>M. guianensis</i>	Tree	S(ls), S, C, GF, MF	63
Myrtaceae	<i>Blepharocalyx salicifolius</i>	Tree	S, C, GF	15
	<i>Campomanesia pubescens</i>	Shrub	S, GF	248
	<i>Eugenia bracteata</i>	Shrub	S(ls), S, GF	174
	<i>E. dysenterica</i>	Tree	S, C, GF	--
	<i>E. puniceifolia</i>	Shrub	S, C, MF	451
	<i>Myrcia cordifolia</i>	Shrub	S, F	45
	<i>M. nivea</i>	Shrub	S(ls), S, C, GF, MF	65
	<i>M. rostrata</i>	Tree	S	105
	<i>M. rubella</i>	Subshrub	S(ls), S	27
	<i>M. tomentosa</i>	Tree	S, C, GF, MF	100
	<i>Pimenta pseudocaryophyllus</i>	Tree	S, GF	446
	<i>Psidium firmum</i>	Shrub	S, GF	595
	<i>P. laruotteanum</i>	Shrub	S, C, GF	157
	<i>Siphoneugena densiflora</i>	Tree	S, C, GF	17
Nyctaginaceae	<i>Guapira areolata</i>	Tree	GF	--
	<i>G. graciliflora</i>	Tree	S(ls), S, C, GF	12
	<i>G. noxia</i>	Tree	S, C, GF	602
	<i>G. opposita</i>	Tree	S(ls), GF, MF	23
	<i>Neea theifera</i>	Tree	S, C, GF	161
Ochnaceae	<i>Ouratea confertiflora</i>	Shrub	S, C	121
	<i>O. floribunda</i>	Subshrub	S, GF	474
	<i>O. hexasperma</i>	Shrub	S, C	40
	<i>O. riedeliana</i>	Shrub	S	180
Opiliaceae	<i>Agonandra brasiliensis</i>	Tree	S, C, GF, MF	--
Orchidaceae	<i>Oncidium varicosum</i>	Herb	GF	318
Oxalidaceae	<i>Oxalis suborbicularis</i>	Subshrub	S(ls), S, C, GF	206
Passifloraceae	<i>Passiflora cerradensis</i>	Climber	C, GF	610
Peraceae	<i>Pera glabrata</i>	Shrub	S(ls), S, C, GF, MF	38
Piperaceae	<i>Piper aduncum</i>	Shrub	C, GF	273
Poaceae	<i>Aristida pendula</i>	Herb	S(ls)	523
	<i>Axonopus barbigerus</i>	Herb	S	524
	<i>Echinolaena inflexa</i>	Herb	S	339
	<i>Ichnanthus pallens</i>	Herb	S, GF	288
	<i>I. bambusiflorus</i>	Herb	S, F, GF	616
	<i>Melinis minutiflora</i> *	Herb	S, GF	287
	<i>Mesosetum loliforme</i>	Herb	S	620
	<i>Panicum cervicatum</i>	Herb	S, C	367

continue

continuation

Family	Species	Stratum	Phytophysiognomy	Coll. N.
	<i>P. maximum</i> *	Herb	S(ls), GF	298
	<i>P. selowii</i>	Herb	S, C, F, GF	410
	<i>Paspalum polyphyllum</i>	Herb	S, GF	521
	<i>Schizachyrium sanguineum</i> *	Herb	S, GF	619
	<i>S. tenerum</i>	Herb	S, C	621
	<i>Trachypogon</i> sp.1	Herb	--	623
	<i>Trachypogon</i> sp.2	Herb	--	624
	<i>Urochloa brizantha</i> *	Herb	S(ls), GF	615
	<i>U. decumbens</i> *	Herb	S	264
Polygalaceae	<i>Bredemeyera velutina</i>	Shrub	S(ls), S, GF	295
	<i>Polygala violacea</i>	Herb	S	257
Proteaceae	<i>Roupala montana</i>	Tree	S, GF	239
Rubiaceae	<i>Alibertia elliptica</i>	Shrub	S(ls), S, F, GF	80
	<i>A. macrophylla</i>	Tree	S(ls), S, C, GF, MF	--
	<i>A. sessilis</i>	Tree	S(ls), S, GF, MF	144
	<i>Amaioua guianensis</i>	Tree	S(ls), GF	--
	<i>Chomelia ribesoides</i>	Subshrub	S, GF	234
	<i>Coccocypselum aureum</i>	Herb	S(ls), S, C, GF	398
	<i>Declieuxia fruticosa</i>	Subshrub	S, GF	382
	<i>Ferdinandusa elliptica</i>	Tree	S, GF	18
	<i>Galium noxium</i>	Herb	GF	265
	<i>Palicourea marcgravii</i>	Subshrub	C, GF	284
	<i>P. officinalis</i>	Subshrub	S, GF	279
	<i>P. rigida</i>	Shrub	S, GF	241
	<i>Psychotria hoffmannseggiana</i>	Subshrub	C, GF	297
	<i>P. prunifolia</i>	Subshrub	C, GF	272
	<i>P. sciaphila</i>	Subshrub	S(ls), C, GF	291
	<i>Sabicea brasiliensis</i>	Subshrub	S, GF	334
	<i>Spermacoce poaya</i>	Herb	S(ls), S	374
	<i>Tocoyena formosa</i>	Shrub	S, C, GF	245
Rutaceae	<i>Esenbeckia pumila</i>	Shrub	S, GF	208
	<i>Spiranthera odoratissima</i>	Shrub	S, C, GF	575
	<i>Zanthoxylum rhoifolium</i>	Tree	S(ls), C, GF	606
Salicaceae	<i>Casearia grandiflora</i>	Shrub	S(ls), S, C, GF, MF	346
	<i>C. sylvestris</i>	Shrub	S, C, GF	552
Sapindaceae	<i>Matayba guianensis</i>	Tree	S, C, GF, MF	220
	<i>Serjania erecta</i>	Climber	S, F, GF	375
	<i>S. lethalis</i>	Climber	S(ls), S, GF	160
	<i>S. ovalifolia</i>	Climber	S(ls), GF	111
Sapotaceae	<i>Pouteria ramiflora</i>	Tree	S, C, GF	576
Schizaeaceae	<i>Anemia phyllitidis</i>	Herb	S, GF, MF	269
Simaroubaceae	<i>Simarouba versicolor</i>	Tree	S, C, GF, MF	108
Siparunaceae	<i>Siparuna guianensis</i>	Tree	S, C, GF, MF	261
Smilacaceae	<i>Smilax brasiliensis</i>	Climber	S, MF	190
Solanaceae	<i>Solanum paniculatum</i> *	Shrub	S, C, GF	116
Styracaceae	<i>Styrax camporum</i>	Tree	S, C, GF, MF	251
	<i>S. ferrugineus</i>	Tree	S, C, GF	39
Symplocaceae	<i>Symplocos mosenii</i>	Tree	S(ls), S, C, GF	11
	<i>S. nitens</i>	Tree	F, GF	455
Turneraceae	<i>Turnera lamiifolia</i>	Herb	S(ls), S	488
Verbenaceae	<i>Lantana camara</i> *	Shrub	S(ls), GF, MF	296
	<i>L. trifolia</i> *	Shrub	S, MF	266

continue

continuation

Family	Species	Stratum	Phytophysiognomy	Coll. N.
Vochysiaceae	<i>Lippia rotundifolia</i>	Shrub	S	511
	<i>Stachytarpheta polyura</i>	Herb	S(ls), S, GF	293
	<i>Callisthene major</i>	Tree	S(ls), S, GF, MF	596
	<i>Qualea dichotoma</i>	Tree	S(ls), S, C, GF, MF	--
	<i>Q. grandiflora</i>	Tree	S, C, GF	283
	<i>Q. multiflora</i>	Tree	S, C, GF	104
	<i>Q. parviflora</i>	Tree	S, C, GF	56
	<i>Vochysia elliptica</i>	Tree	S	--
	<i>V. rufa</i>	Tree	S, C, GF	--
	<i>V. thyrsoides</i>	Tree	S, C	165
	<i>V. tucanorum</i>	Tree	S(ls), S, C, GF, MF	49

Higher proportions of arboreal species in cerradão sites as compared to herbaceous or shrub forms are not typical of the Cerrado biome as a whole, which consistently shows higher proportions of species in the herbaceous stratum. Mendonça et al. (2008) noted, however, that all of the subshrubs, climbers and lianas they identified as belonging to the herbaceous stratum could be considered as belonging to the shrub or even arboreal layer (as they were reasonably large, as in the case of woody vines). Other lists, including that obtained in the present study, have confirmed the predominance of the arboreal stratum in the cerradão phytobiognomies.

Batalha & Mantovani (2001) reported 148 species belonging to 113 genera and 45 families in an area of cerradão in São Paulo State, Brazil, with a proportion of herbaceous to woody species (0.8:1.0) similar to that of the present study. Comparisons of the richness values between any two areas, however, must take into consideration the collection frequencies and the flora types considered (vascular, or only phanerogams).

The families Fabaceae, Asteraceae, Rubiaceae, Poaceae, Malpighiaceae, Myrtaceae, Melastomataceae, Apocynaceae, and Vochysiaceae had the greatest species richness in the present study (52.12% of the total number of species encountered). In relation to the numbers of genera, Fabaceae was again the most abundant family, followed by Asteraceae, Rubiaceae, Poaceae, Myrtaceae, and Bignoniaceae; *Miconia* was the most representative genus (nine species).

Rubiaceae, Myrtaceae, Fabaceae, Vochysiaceae, and Melastomataceae were also among the families demonstrating the greatest richnesses of woody species in areas of cerradão studied by Pereira-Silva et al. (2004); Fabaceae, Malvaceae, and Myrtaceae stood out among the cerradão families in Pantanal de Nhecolândia in Mato Grosso do Sul State (MS) (Salis et al. 2006); and Fabaceae

and Vochysiaceae were the richest families identified in cerrado sensu stricto sites by Assunção & Felfili (2004) and by Costa & Araújo (2001), with Myrtaceae also showing high richness in the latter survey. Studies undertaken in forest areas demonstrated different results, however. The families Fabaceae, Melastomataceae, Myrtaceae, Rubiaceae, Apocynaceae, and Vochysiaceae were cited as demonstrating the greatest species richness in gallery forests, with important contributions also by Sapotaceae, Euphorbiaceae, and Moraceae (poorly represented in the present study) (Marimon et al. 2002, Silva Junior 2004); and Fabaceae, Bignoniaceae, Malvaceae, and Anacardiaceae were well represented in a semi-deciduous dry forest in Diorama, Goiás State (GO) (Santos-Diniz & Souza 2011). It can therefore be seen that, in spite of the floristic differences between the phytobiognomies, the richness values observed in the Cerradão REBIO were close to those reported for cerrado sensu stricto (Felfili et al. 1994, Costa & Araújo 2001, Silva et al. 2002). According to Souza et al. (2010), the importance of Fabaceae in terms of species richness in the Cerrado biome is due to the capacity of many of its species to fix nitrogen, enabling them to grow in poor and degraded soils.

Lopes (1984) noted that plant distributions are controlled by very basic factors such as the climate, soils, and the availability of water and nutrients, but can also be greatly influenced by latitude, burning frequencies, water table depth, animal pasturing, and anthropogenic factors – which reinforces the importance of floristic and phytosociological studies at many different sites, as each area would be expected to have specific physical conditions and therefore unique population structures (Santos-Diniz & Souza 2011).

The data published in the three plant lists used for comparisons in the present work, in the flora of the

Federal District, and in floristic and phytosociological studies undertaken in both savanna and cerrado forest phytophysiognomies (Felfili et al. 1994, Silva Junior et al. 1998, Marimon et al. 2002, Silva et al. 2002, Assunção & Felfili 2004, Pereira-Silva et al. 2004, Silva Junior 2004, Balduino et al. 2005, Santiago et al. 2005, Salis et al. 2006) indicated that 91% of the fully-identified species encountered in the REBIO cerradão were common to cerrado areas, while 79.5% were common to forest sites. The species commonly encountered in both formations represent 72.3% of the total, while only 60.43% of the fully identified species of the REBIO were reported in other cerradão surveys. No single species was exclusive to the cerradão phytophysiognomy.

The elevated species richness identified in the study area (with almost 95% species being native to the Cerrado biome) indicates the importance of this conservation area, for it not only protects a significant area of cerradão vegetation (a phytophysiognomy poorly represented in other designated conservation areas) but also conserves species such as *Copaifera langsdorffii* (*copaíba*), *Pterodon pubescens* (*sucupira-branca*), *Caryocar brasiliense* (*pequizeiro*), *Eugenia dysenterica* (*cagaiteira*), *Vochysia thyrsoides* (*gomeira*), *Vochysia tucanorum* (*pau-doce*), *Pseudobombax longiflorum* (*embiruçu*), *Dalbergia miscolobium* (*jacarandá*), *Aspidosperma* spp. (*perobas*), and *Tabebuia* spp. (*ipês*) that have been designated as protected elements of the Ecological Heritage of the Federal District (Felfili & Santos 2002).

The results obtained in the present study enrich our knowledge of the cerradão flora and can serve as subsidies for the elaboration of management programs for this conservation area (REBIO).

REFERENCES

- APG II. 2003. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Botanical Journal of the Linnean Society* 141:399-436.
- Assis MC. 2002. Alstroemeriaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.2, p.19-32.
- Assunção SL, Felfili JM. 2004. Fitossociologia de um fragmento de cerrado *sensu stricto* na APA do Paranoá, DF, Brasil. *Acta Botanica Brasilica* 18:903-909.
- Balduino APC, Souza AL, Meira Neto JAA, Silva AF, Silva Junior MC. 2005. Fitossociologia e análise comparativa da composição florística do cerrado da flora de Paraopeba-MG. *Revista Árvore* 29:25-34.
- Batalha MA, Mantovani W. 2001. Floristic composition of the Cerrado in the Pé-de-Gigante Reserve (Santa Rita do Passa Quatro, Southeastern, Brazil). *Acta Botanica Brasilica* 15:289-304.
- Carvalho-Silva M, Cavalcanti TB. 2002. Piperaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.2, p.91-124.
- Chacon RG, Yamamoto K, Cavalcanti TB. 2003. Ochnaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.3, p.205-226.
- Costa AA, Araújo GM. 2001. Comparação da vegetação arbórea de cerradão e de cerrado na Reserva do Panga, Uberlândia, Minas Gerais. *Acta Botanica Brasilica* 15:63-72.
- Dittrich VAO, Heringer G, Salino A. 2007. Blechnaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, org.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.6, p.89-108.
- Felfili JM. 2002. Padrões de diversidade do Cerrado do Centro-Oeste brasileiro. In Biodiversidade, conservação e uso sustentável da flora do Brasil (EL Araújo, AN Moura, ESB Sampaio, LMS Gestinari, JMT Carneiro, eds.). Universidade Federal Rural de Pernambuco – Imprensa Universitária, Recife, p.58-61.
- Felfili JM, Santos AAB. 2002. Legislação Ambiental: APA Gama e Cabeça de Veado. Universidade de Brasília, Faculdade de Tecnologia, Departamento de Engenharia Florestal, Brasília.
- Felfili JM, Haridasan M, Mendonça RC, Filgueiras TS, Silva Junior MC, Rezende AV. 1994. Projeto biogeografia do Bioma Cerrado: vegetação e solos. *Caderno de Geociências* 12:75-166.
- Felfili JM, Mendonça RC, Munhoz CBR, Fagg CW, Pinto JRR, Silva Junior MC, Sampaio JC. 2004. Vegetação e flora da APA Gama e Cabeça de Veado. In Flora e diretrizes ao plano de manejo da APA Gama e Cabeça de Veado (JM Felfili, AAB Santos, JC Sampaio, orgs.). Universidade de Brasília, Faculdade de Tecnologia, Departamento de Engenharia Florestal, Brasília, p.7-16.
- Ferreira JN, Aquino FG, Okano RC, Proença CEB. 2003. Celastraceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.3, p.139-150.
- Fiaschi P. 2006. Oxalidaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, org.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.5, p.91-106.
- França F. 2005. Vochysiaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.4, p.77-106.

- Groppi M. 2003. Aquifoliaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.3, p.49-62.
- Groppi M, Pirani JR. 2007. Rutaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, org.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.6, p.63-86.
- Ibram. 2010. Proposta de mudança de categoria: Área de Relevante Interesse Ecológico do Cerradão – Reserva Biológica do Cerradão. Governo do Distrito Federal, Brasília.
- Lima LR, Pirani JR. 2003. Burseraceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.3, p.125-138.
- Lombardi JA. 2002. Hippocrateaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.2, p.73-90.
- Lopes AS. 1984. Solos sob “cerrado”: características, propriedades e manejo. 2nd ed. Associação Brasileira para Pesquisa da Potassa e do Fosfato, Piracicaba.
- Marimon BS, Felfili JM, Lima ES. 2002. Floristics and phytosociology of the gallery forest of the Bacaba Stream, Nova Xavantina, Mato Grosso, Brazil. Edinburgh Journal of Botany 59: 303-318.
- Martins RC, Filgueiras TS. 2006. Arecaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, org.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.5, p.45-82.
- Mendonça JO, Amaral Junior A. 2002. Erythroxylaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.2, p.53-71.
- Mendonça RC, Felfili JM, Walter BMT, Silva Junior MC, Rezende AB, Filgueiras TS, Nogueira PE, Fagg CW. 2008. Flora vascular do bioma Cerrado: checklist com 12.356 espécies. In Cerrado: ecologia e flora (SM Sano, SP Almeida, JF Ribeiro, eds.). Embrapa Cerrados/Embrapa Informação Tecnológica, Brasília, v.2, p.421-1279.
- Missouri Botanical Garden – W³Tropicos. 2008. <http://www.tropicos.org> (accessed 2008 Nov 07).
- Pastore JFB, Cavalcanti TB. 2006. Polygalaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, org.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.5, p.107-180.
- Pereira-Silva EFL, Santos JE, Kageyama PY, Hardt E. 2004. Florística e fitossociologia dos estratos arbustivo e arbóreo de um remanescente de cerradão em uma unidade de conservação do Estado de São Paulo. Revista Brasileira de Botânica 27: 533-544.
- Pontes AF, Mello-Silva R. 2003. Annonaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.3, p.19-48.
- Proença CEB, Munhoz CBR, Jorge CL, Nóbrega MGG. 2001. Listagem e nível de proteção das espécies de fanerógamas do Distrito Federal, Brasil. In Flora do Distrito Federal I (TB Cavalcanti, AE Ramos, eds.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, p.89-359.
- Ribeiro JF, Walter BMT. 2008. As principais fitofisionomias do bioma Cerrado. In Cerrado: ecologia e flora (SM Sano, SP Almeida, JF Ribeiro, eds.). Embrapa Cerrados/Embrapa Informação Tecnológica, Brasília, 2nd ed., p.151-212.
- Ribeiro MF, Proença CEB. 2002. Caryocaraceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.2, p.33-38.
- Salis SM, Assis M, Crispim SMA, Casagrande JC. 2006. Distribuição e abundância de espécies arbóreas em cerradões no Pantanal, Estado do Mato Grosso do Sul, Brasil. Revista Brasileira de Botânica 29:339-352.
- Santiago J, Silva Junior MC, Lima LC. 2005. Fitossociologia da regeneração arbórea na mata de galeria do Pitoco (IBGE-DF), seis anos após fogo acidental. Scientia Forestalis 67:64-77.
- Santos-Diniz VS, Sousa TD. 2011. Levantamento florístico e fitossociológico de mata seca semidecídua em área de Reserva Legal do município de Diorama, região oeste de Goiás, Brasil. Encyclopédia Biosfera 7:1-17.
- Silva JSS. 2009. Diversidade alfa, florística e fitossociologia na ARIE do Cerradão, na APA Gama e Cabeça de Veado, DF. Dissertação de mestrado, Universidade de Brasília, Brasília.
- Silva LO, Costa DA, Espírito-Santo Filho K, Ferreira HD, Brandão D. 2002. Levantamento florístico e fitossociológico em duas áreas de cerrado *sensu stricto* no Parque Estadual da Serra de Caldas Novas, Goiás. Acta Botanica Brasiliensis 16:43-53.
- Silva MRP, Barros ICL. 2005. Schizaeaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.4, p.217-248.
- Silva NMF, Valente MC. 2003. Combretaceae. In Flora do Distrito Federal, Brasil (TB Cavalcanti, AE Ramos, orgs.). Embrapa Recursos Genéticos e Biotecnologia, Brasília, v.3, p.151-170.
- Silva Junior MC. 2004. Fitossociologia e estrutura diamétrica da Mata de galeria do Taquara, na Reserva Ecológica do IBGE, DF. Revista Árvore 28:419-428.
- Silva Junior MC, Felfili JM, Nogueira PE, Rezende AB. 1998. Análise florística das matas de galeria no Distrito Federal. In Cerrado: matas de galeria (JF Ribeiro, ed.). Embrapa Cerrados, Planaltina, p.53-84.

- Smith AR, Pryer KM, Schuettpelz E, Korall P, Schneider H, Wolf PG. 2006. A classification for extant ferns. *Taxon* 55:705-731.
- Souza PB, Saporetti Junior AW, Soares MP, Viana RHO, Camargos VL, Meira Neto JAA. 2010. Florística de uma área de cerradão na Floresta Nacional de Paraopeba – Minas Gerais. *Cerne* 16:86-93.
- The International Plant Names Index – IPNI. 2008. <http://www.ipni.org> (accessed 2008 Nov 07).
- Unesco. 2002. Vegetação do Distrito Federal: tempo e espaço – Uma avaliação multitemporal da perda de cobertura vegetal no DF e da diversidade florística da Reserva da Biosfera do Cerrado – Fase I. 2nd ed., Unesco, Brasília.

